THE FIRST SIX BOOKS OF EUCLID.



WITH

COLOURED DIAGRAMS

BY

BYRNE

Harvard University - Houghton Library / pga_typ_805_47_3730

Philip Hofer

Presented by Mr M Dick
To his young friend
Mr Robt Gardner Jr.
Jany 1858.

HARVARD COLLEGE LIBRARY
From the collection of Philip Hofer
Class of 1921 for the DEPARTMENT
OF PRINTING AND GRAPHIC ARTS.

Harvard University - Houghton Library / pga_typ_805_47_3730

Harvard University - Houghton Library / pga_typ_805_47_3730

Scarce

£3

Harvard University - Houghton Library / pga_typ_805_47_3730

BYRNE'S EUCLID

THE FIRST SIX BOOKS OF

THE ELEMENTS OF EUCLID

WITH COLOURED DIAGRAMS

AND SYMBOLS

Harvard University - Houghton Library / pga_typ_805_47_3730

Harvard University - Houghton Library / pga_typ_805_47_3730

THE FIRST SIX BOOKS OF

THE ELEMENTS OF EUCLID

IN WHICH COLOURED DIAGRAMS AND SYMBOLS


ARE USED INSTEAD OF LETTERS FOR THE

GREATER EASE OF LEARNERS

BY OLIVER BYRNE

SURVEYOR OF HER MAJESTY'S SETTLEMENTS IN THE FALKLAND ISLANDS

AND AUTHOR OF NUMEROUS MATHEMATICAL WORKS



LONDON

WILLIAM PICKERING

1847

Harvard University - Houghton Library / pga_typ_805_47_3730

Harvard University - Houghton Library / pga_typ_805_47_3730

TO THE

RIGHT HONOURABLE THE EARL FITZWILLIAM,

ETC. ETC. ETC.

THIS WORK IS DEDICATED

BY HIS LORDSHIP'S OBEDIENT

AND MUCH OBLIGED SERVANT,

OLIVER BYRNE.

Harvard University - Houghton Library / pga_typ_805_47_3730

Harvard University - Houghton Library / pga_typ_805_47_3730

INTRODUCTION.

THE arts and ſciences have become ſo extenſive, that to facilitate their acquirement is of as much importance as to extend their boundaries. Illuſtration, if it does not ſhorten the time of ſtudy, will at leaſt make it more agreeable. THIS WORK has a greater aim than mere illuſtration; we do not introduce colours for the purpoſe of entertainment, or to amuſe *by certain combinations of tint and form*, but to aſſiſt the mind in its reſearches after truth, to increaſe the facilities of inſtruction, and to diffuſe permanent knowledge. If we wanted authorities to prove the importance and uſefulneſs of geometry, we might quote every philoſopher ſince the days of Plato. Among the Greeks, in ancient, as in the ſchool of Peſtalozzi and others in recent times, geometry was adopted as the beſt gymnaſtic of the mind. In fact, Euclid's Elements have become, by common conſent, the baſis of mathematical ſcience all over the civilized globe. But this will not appear extraordinary, if we conſider that this ſublime ſcience is not only better calculated than any other to call forth the ſpirit of inquiry, to elevate the mind, and to ſtrengthen the reaſoning faculties, but alſo it forms the beſt introduction to moſt of the uſeful and important vocations of human life. Arithmetic, land-ſurveying, menſuration, engineering, navigation, mechanics, hydroſtatics, pneumatics, optics, phyſical aſtronomy, &c. are all dependent on the propoſitions of geometry.

Harvard University - Houghton Library / pga_typ_805_47_3730

Much however depends on the firſt communication of any ſcience to a learner, though the beſt and moſt eaſy methods are ſeldom adopted. Propoſitions are placed before a ſtudent, who though having a ſufficient underſtanding, is told juſt as much about them on entering at the very threſhold of the ſcience, as gives him a prepoſſeſſion moſt unfavourable to his future ſtudy of this delightful ſubject; or "the formalities and paraphernalia of rigour are ſo oſtentatiouſly put forward, as almoſt to hide the reality. Endleſs and perplexing repetitions, which do not confer greater exactitude on the reaſoning, render the demonſtrations involved and obſcure, and conceal from the view of the ſtudent the conſecution of evidence." Thus an averſion is created in the mind of the pupil, and a ſubject ſo calculated to improve the reaſoning powers, and give the habit of cloſe thinking, is degraded by a dry and rigid courſe of inſtruction into an unintereſting exerciſe of the memory. To raiſe the curioſity, and to awaken the liſtleſs and dormant powers of younger minds ſhould be the aim of every teacher; but where examples of excellence are wanting, the attempts to attain it are but few, while eminence excites attention and produces imitation. The object of this Work is to introduce a method of teaching geometry, which has been much approved of by many ſcientific men in this country, as well as in France and America. The plan here adopted forcibly appeals to the eye, the moſt ſenſitive and the moſt comprehenſive of our external organs, and its pre-eminence to imprint it ſubject on the mind is ſupported by the incontrovertible maxim expreſſed in the well known words of Horace:—


Segnius irritant animos demiſſa per aurem
Quàm quæ ſunt oculis ſubjecta fidelibus.

A feebler impreſs through the ear is made,
Than what is by the faithful eye conveyed.

Harvard University - Houghton Library / pga_typ_805_47_3730

All language conſiſts of repreſentative ſigns, and thoſe ſigns are the beſt which effect their purpoſes with the greateſt preciſion and diſpatch. Such for all common purpoſes are the audible ſigns called words, which are ſtill conſidered as audible, whether addreſſed immediately to the ear, or through the medium of letters to the eye. Geometrical diagrams are not ſigns, but the materials of geometrical ſcience, the object of which is to ſhow the relative quantities of their parts by a proceſs of reaſoning called Demonſtration. This reaſoning has been generally carried on by words, letters, and black or uncoloured diagrams; but as the uſe of coloured ſymbols, ſigns, and diagrams in the linear arts and ſciences, renders the proceſs of reaſoning more preciſe, and the attainment more expeditious, they have been in this inſtance accordingly adopted.

Such is the expedition of this enticing mode of communicating knowledge, that the Elements of Euclid can be acquired in leſs than one third the time uſually employed, and the retention by the memory is much more permanent; theſe facts have been aſcertained by numerous experiments made by the inventor, and ſeveral others who have adopted his plans. The particulars of which are few and obvious; the letters annexed to points, lines, or other parts of a diagram are in fact but arbitrary names, and repreſent them in the demonſtration; inſtead of theſe, the parts being differently coloured, are made to name themſelves, for their forms in correſponding colours represent them in the demonſtration.



In order to give a better idea of this ſyſtem, and of the advantages gained by its adoption, let us take a right

Harvard University - Houghton Library / pga_typ_805_47_3730

angled triangle, and expreſs ſome of its properties both by colours and the method generally employed.

Some of the properties of the right angled triangle ABC, *expreſſed by the method generally employed.*

1. The angle BAC, together with the angles BCA and ABC are equal to two right angles, or twice the angle ABC.

2. The angle CAB added to the angle ACB will be equal to the angle ABC.


3. The angle ABC is greater than either of the angles BAC or BCA.

4. The angle BCA or the angle CAB is leſs than the angle ABC.

5. If from the angle ABC, there be taken the angle BAC, the remainder will be equal to the angle ACB.

6. The ſquare of AC is equal to the ſum of the ſquares of AB and BC.

The ſame properties expreſſed by colouring the different parts.

1. 

That is, the red angle added to the yellow angle added to the blue angle, equal twice the yellow angle, equal two right angles.

2. 

Or in words, the red angle added to the blue angle, equal the yellow angle.


3. 

The yellow angle is greater than either the red or blue angle.


Harvard University - Houghton Library / pga_typ_805_47_3730

4.  .

Either the red or blue angle is leſs than the yellow angle.

5.  .

In other terms, the yellow angle made leſs by the blue angle equal the red angle.

6.  .

That is, the ſquare of the yellow line is equal to the ſum of the ſquares of the blue and red lines.

In oral demonſtrations we gain with colours this important advantage, the eye and the ear can be addreſſed at the ſame moment, ſo that for teaching geometry, and other linear arts and ſciences, in claſſes, the ſyſtem is the beſt ever propoſed, this is apparent from the examples juſt given.

Whence it is evident that a reference from the text to the diagram is more rapid and ſure, by giving the forms and colours of the parts, or by naming the parts and their colours, than naming the parts and letters on the diagram. Beſides the ſuperior ſimplicity, this ſyſtem is likewiſe conſpicuous for concentration, and wholly excludes the injurious though prevalent practice of allowing the ſtudent to commit the demonſtration to memory; until reaſon, and fact, and proof only make impreſſions on the underſtanding.

Again, when lecturing on the principles or properties of figures, if we mention the colour of the part or parts referred to, as in ſaying, the red angle, the blue line, or lines, &c. the part or parts thus named will be immediately ſeen by all in the claſs at the ſame inſtant; not ſo if we ſay the angle ABC, the triangle PFQ, the figure EGKt, and ſo on;

Harvard University - Houghton Library / pga_typ_805_47_3730

tion of three colours on a plane, gives a good idea of a mathematical point.


A line is length without breadth. With the aſſiſtance of colours, nearly in the ſame manner as before, an idea of a line may be thus given:—

Let two colours meet and cover a portion of the paper;



where they meet is not red, nor is it blue; therefore the junction occupies no portion of the plane, and therefore it cannot have breadth, but only length: from which we can readily form an idea of what is meant by a mathematical line. For the purpoſe of illuſtration, one colour differing from the colour of the paper, or plane upon which it is drawn, would have been ſufficient; hence in future, if we ſay the red line, the blue line, or lines, &c. it is the junctions with the plane upon which they are drawn are to be underſtood.

Surface is that which has length and breadth without thickneſs.




When we conſider a ſolid body (PQ), we perceive at once that it has three dimenſions, namely:—length, breadth, and thickneſs; ſuppoſe one part of this ſolid (PS) to be red, and the other part (QR) yellow, and that the colours be diſtinct without commingling, the blue ſurface (RS) which ſeparates theſe parts, or which is the ſame thing, that which divides the ſolid without loſs of material, muſt be without thickneſs, and only poſſeſſes length and breadth;

this plainly appears from reaſoning, ſimilar to that juſt employed in defining, or rather deſcribing a point and a line.

The propoſition which we have ſelected to elucidate the manner in which the principles are applied, is the fifth of the firſt Book.

In an iſoſceles triangle ABC, the internal angles at the baſe ABC, ACB are equal, and when the ſides AB, AC are produced, the external angles at the baſe BCE, CBD are alſo equal.




Produce ——— and ———
make ——— = ———
Draw ——— and ———
(B. 1. pr. 3.)

in  and 

we have ——— ——— = ——— ———

——— = ——— and  common :




∴ ——— = ———,

and  =  (B. 1. pr. 4.)

Again in  and  ,

——— = ———,

Harvard University - Houghton Library / pga_typ_805_47_3730



Q. E. D.

By annexing Letters to the Diagram.

Let the equal ſides AB and AC be produced through the extremities BC, of the third ſide, and in the produced part BD of either, let any point D be aſſumed, and from the other let AE be cut off equal to AD (B. 1. pr. 3). Let the points E and D, ſo taken in the produced ſides, be connected by ſtraight lines DC and BE with the alternate extremities of the third ſide of the triangle.

In the triangles DAC and EAB the ſides DA and AC are reſpectively equal to EA and AB, and the included angle A is common to both triangles. Hence (B. 1. pr. 4.) the line DC is equal to BE, the angle ADC to the angle AEB, and the angle ACD to the angle ABE; if from the equal lines AD and AE the equal ſides AB and AC be taken, the remainders BD and CE will be equal. Hence in the triangles BDC and CEB, the ſides BD and DC are reſpectively equal to CE and EB, and the angles D and E included by thoſe ſides are alſo equal. Hence (B. 1. pr. 4.)

Harvard University - Houghton Library / pga_typ_805_47_3730

the angles DBC and ECB, which are thoſe included by the third ſide BC and the productions of the equal ſides AB and AC are equal. Alſo the angles DCB and EBC are equal if thoſe equals be taken from the angles DCA and EBA before proved equal, the remainders, which are the angles ABC and ACB oppoſite to the equal ſides, will be equal.

Therefore in an iſoſceles triangle, &c.

Q. E. D.

Our object in this place being to introduce the ſyſtem rather than to teach any particular ſet of propoſitions, we have therefore ſelected the foregoing out of the regular courſe. For ſchools and other public places of inſtruction, dyed chalks will anſwer to deſcribe diagrams, &c. for private uſe coloured pencils will be found very convenient.

We are happy to find that the Elements of Mathematics now forms a conſiderable part of every ſound female education, therefore we call the attention of thoſe intereſted or engaged in the education of ladies to this very attractive mode of communicating knowledge, and to the ſucceeding work for its future developement.

We ſhall for the preſent conclude by obſerving, as the ſenſes of ſight and hearing can be ſo forcibly and inſtantaneously addreſſed alike with one thouſand as with one, *the million* might be taught geometry and other branches of mathematics with great eaſe, this would advance the purpoſe of education more than any thing that *might* be named, for it would teach the people how to think, and not what to think; it is in this particular the great error of education originates.

Harvard University - Houghton Library / pga_typ_805_47_3730

THE ELEMENTS OF EUCLID.

BOOK I.

DEFINITIONS.

I.

A *point* is that which has no parts.

II.

A *line* is length without breadth.

III.

The extremities of a line are points.

IV.

A ſtraight or right line is that which lies evenly between its extremities.

V.

A ſurface is that which has length and breadth only.

VI.

The extremities of a ſurface are lines.


VII.

A plane ſurface is that which lies evenly between its extremities.

VIII.

A plane angle is the inclination of two lines to one another, in a plane, which meet together, but are not in the ſame direction.

IX.



A plane rectilinear angle is the inclination of two ſtraight lines to one another, which meet together, but are not in the ſame ſtraight line.

Harvard University - Houghton Library / pga_typ_805_47_3730

X.

When one ſtraight line ſtanding on another ſtraight line makes the adjacent angles equal, each of theſe angles is called a *right angle*, and each of theſe lines is ſaid to be *perpendicular* to the other.



XI.

An obtuſe angle is an angle greater than a right angle.



XII.

An acute angle is an angle leſs than a right angle.



XIII.


A term or boundary is the extremity of any thing.

XIV.

A figure is a ſurface encloſed on all ſides by a line or lines.

XV.

A circle is a plane figure, bounded by one continued line, called its circumference or periphery; and having a certain point within it, from which all ſtraight lines drawn to its circumference are equal.



XVI.

This point (from which the equal lines are drawn) is called the centre of the circle.

Harvard University - Houghton Library / pga_typ_805_47_3730




XVII.

A diameter of a circle is a ſtraight line drawn through the centre, terminated both ways in the circumference.



XVIII.

A ſemicircle is the figure contained by the diameter, and the part of the circle cut off by the diameter.



XIX.

A ſegment of a circle is a figure contained by a ſtraight line, and the part of the circumference which it cuts off.


XX.

A figure contained by ſtraight lines only, is called a rectilinear figure.

XXI.

A triangle is a rectilinear figure included by three ſides.

XXII.



A quadrilateral figure is one which is bounded by four ſides. The ſtraight lines —— and —— connecting the vertices of the oppoſite angles of a quadrilateral figure, are called its diagonals.

XXIII.

A polygon is a rectilinear figure bounded by more than four ſides.

Harvard University - Houghton Library / pga_typ_805_47_3730

XXIV.

A triangle whoſe three ſides are equal, is ſaid to be equilateral.



XXV.

A triangle which has only two ſides equal is called an iſoſceles triangle.



XXVI.

A ſcalene triangle is one which has no two ſides equal.


XXVII.

A right angled triangle is that which has a right angle.



XXVIII.

An obtuſe angled triangle is that which has an obtuſe angle.



XXIX.

An acute angled triangle is that which has three acute angles.




XXX.

Of four-ſided figures, a ſquare is that which has all its ſides equal, and all its angles right angles.




XXXI.

A rhombus is that which has all its ſides equal, but its angles are not right angles.




XXXII.

An oblong is that which has all its angles right angles, but has not all its ſides equal.



Harvard University - Houghton Library / pga_typ_805_47_3730

XXXIII.




A rhomboid is that which has its oppoſite ſides equal to one another, but all its ſides are not equal, nor its angles right angles.

XXXIV.

All other quadrilateral figures are called trapeziums.

XXXV.



Parallel ſtraight lines are ſuch as are in the ſame plane, and which being produced continually in both directions, would never meet.

POSTULATES.

I.

Let it be granted that a ſtraight line may be drawn from any one point to any other point.

II.

Let it be granted that a finite ſtraight line may be produced to any length in a ſtraight line.

III.

Let it be granted that a circle may be deſcribed with any centre at any diſtance from that centre.

AXIOMS.

I.

Magnitudes which are equal to the ſame are equal to each other.

II.

If equals be added to equals the ſums will be equal.

Harvard University - Houghton Library / pga_typ_805_47_3730

III.

If equals be taken away from equals the remainders will be equal.

IV.

If equals be added to unequals the ſums will be unequal.

V.

If equals be taken away from unequals the remainders will be unequal.

VI.

The doubles of the ſame or equal magnitudes are equal.

VII.

The halves of the ſame or equal magnitudes are equal.

VIII.

Magnitudes which coincide with one another, or exactly fill the ſame ſpace, are equal.

IX.

The whole is greater than its part.

X.

Two ſtraight lines cannot include a ſpace.

XI.

All right angles are equal.

XII.

If two ſtraight lines ( ) meet a third ſtraight line ( ) ſo as to make the two interior angles (and) on the ſame ſide leſs than two right angles, theſe two ſtraight lines will meet if they be produced on that ſide on which the angles are leſs than two right angles.



Harvard University - Houghton Library / pga_typ_805_47_3730

The twelfth axiom may be expreſſed in any of the following ways:


1. Two diverging ſtraight lines cannot be both parallel to the ſame ſtraight line.

2. If a ſtraight line interſect one of the two parallel ſtraight lines it muſt alſo interſect the other.


3. Only one ſtraight line can be drawn through a given point, parallel to a given ſtraight line.

Geometry has for its principal objects the expoſition and explanation of the properties of *figure*, and figure is defined to be the relation which ſubſiſts between the boundaries of ſpace. Space or magnitude is of three kinds, *linear*, *ſuperficial*, and *ſolid*.

Angles might properly be conſidered as a fourth ſpecies of magnitude. Angular magnitude evidently conſiſts of parts, and muſt therefore be admitted to be a ſpecies of quantity The ſtudent muſt not ſuppoſe that the magnitude of an angle is affected by the length of the ſtraight lines which include it, and of whoſe mutual divergence it is the meaſure. The *vertex* of an angle is the point where the *ſides* or the *legs* of the angle meet, as A.



An angle is often deſignated by a ſingle letter when its legs are the only lines which meet together at its vertex. Thus the red and blue lines form the yellow angle, which in other ſyſtems would be called the angle A. But when more than two lines meet in the ſame point, it was neceſſary by former methods, in order to avoid confuſion, to employ three letters to deſignate an angle about that point,



Harvard University - Houghton Library / pga_typ_805_47_3730

the letter which marked the vertex of the angle being always placed in the middle. Thus the black and red lines meeting together at C, form the blue angle, and has been uſually denominated the angle FCD or DCF. The lines FC and CD are the legs of the angle; the point C is its vertex. In like manner the black angle would be deſignated the angle DCB or BCD. The red and blue angles added together, or the angle HCF added to FCD, make the angle HCD; and ſo of other angles.

When the legs of an angle are produced or prolonged beyond its vertex, the angles made by them on both ſides of the vertex are ſaid to be *vertically oppoſite* to each other: Thus the red and yellow angles are ſaid to be vertically oppoſite angles.

Superpoſition is the proceſs by which one magnitude may be conceived to be placed upon another, ſo as exactly to cover it, or ſo that every part of each ſhall exactly coincide.

A line is ſaid to be *produced*, when it is extended, prolonged, or has its length increaſed, and the increaſe of length which it receives is called its *produced part*, or its *production*.

The entire length of the line or lines which encloſe a figure, is called its *perimeter*. The firſt ſix books of Euclid treat of plain figures only. A line drawn from the centre of a circle to its circumference, is called a *radius*. The lines which include a figure are called its *ſides*. That ſide of a right angled triangle, which is oppoſite to the right angle, is called the *hypotenuſe*. An *oblong* is defined in the ſecond book, and called a *rectangle*. All the lines which are conſidered in the firſt ſix books of the Elements are ſuppoſed to be in the ſame plane.

The *ſtraight-edge* and *compaſſes* are the only inſtruments,

Harvard University - Houghton Library / pga_typ_805_47_3730

the uſe of which is permitted in Euclid, or plain Geometry. To declare this reſtriction is the object of the *poſtulates*.

The *Axioms* of geometry are certain general propoſitions, the truth of which is taken to be ſelf-evident and incapable of being eſtabliſhed by demonſtration.

Propoſitions are thoſe reſults which are obtained in geometry by a proceſs of reaſoning. There are two ſpecies of propoſitions in geometry, *problems* and *theorems*.

A *Problem* is a propoſition in which ſomething is propoſed to be done; as a line to be drawn under ſome given conditions, a circle to be deſcribed, ſome figure to be conſtructed, &c.

The *ſolution* of the problem conſiſts in ſhowing how the thing required may be done by the aid of the rule or ſtraight-edge and compaſſes.

The *demonſtration* conſiſts in proving that the proceſs indicated in the ſolution really attains the required end.

A *Theorem* is a propoſition in which the truth of ſome principle is aſſerted. This principle muſt be deduced from the axioms and definitions, or other truths previously and independently eſtabliſhed. To ſhow this is the object of demonſtration.

A *Problem* is analogous to a poſtulate.

A *Theorem* reſembles an axiom.

A *Poſtulate* is a problem, the ſolution of which is aſſumed.

An *Axiom* is a theorem, the truth of which is granted without demonſtration.

A *Corollary* is an inference deduced immediately from a propoſition.

A *Scholium* is a note or obſervation on a propoſition not containing an inference of ſufficient importance to entitle it to the name of a *corollary*.

A *Lemma* is a propoſition merely introduced for the purpoſe of eſtabliſhing ſome more important propoſition.

Harvard University - Houghton Library / pga_typ_805_47_3730

SYMBOLS AND ABBREVIATIONS.

$\therefore$ expreſſes the word *therefore*.

$\because$ . . . . . *becauſe*.

$=$ . . . . . *equal*. This ſign of equality may be read *equal to*, or *is equal to*, or *are equal to*; but any diſcrepancy in regard to the introduction of the auxiliary verbs *is*, *are*, &c. cannot affect the geometrical rigour.

$\neq$ means the ſame as if the words '*not equal*' were written.

$\sqsubset$ ſignifies *greater than*.

$\sqsupset$ . . . . *leſs than*.

$\not\sqsubset$ . . . . *not greater than*.

$\not\sqsupset$ . . . . *not leſs than*.

$+$ is read *plus* (*more*), the ſign of addition; when interpoſed between two or more magnitudes, ſignifies their ſum.

$-$ is read *minus* (*leſs*), ſignifies ſubtraction; and when placed between two quantities denotes that the latter is to be taken from the former.

$\times$ this ſign expreſſes the product of two or more numbers when placed between them in arithmetic and algebra; but in geometry it is generally uſed to expreſs a *rectangle*, when placed between "two ſtraight lines which contain one of its right angles." A *rectangle* may alſo be repreſented by placing a point between two of its conterminous ſides.

$:\ ::\ :$ expreſſes an *analogy* or *proportion*; thus, if A, B, C and D, repreſent four magnitudes, and A has to B the ſame ratio that C has to D, the propoſition is thus briefly written,

$$A : B :: C : D,$$

$$A : B = C : D,$$

$$\text{or } \frac{A}{B} = \frac{C}{D}.$$


This equality or ſameneſs of ratio is read,


Harvard University - Houghton Library / pga_typ_805_47_3730

as A is to B, ſo is C to D;
or A is to B, as C is to D.


$\parallel$ ſignifies *parallel to.*

$\perp$ . . . . *perpendicular to.*

 . *angle.*

 . . *right angle.*

 *two right angles.*

 briefly deſignates a *point.*

$\sqsubset$, $=$, or $\sqsupset$ ſignifies *greater, equal, or leſs than.*

The ſquare deſcribed on a line is conciſely written thus, $\overline{}^2$.

In the ſame manner twice the ſquare of, is expreſſed by $2 \cdot \overline{}^2$.

def. ſignifies *definition.*

pos. . . . . *poſtulate.*

ax. . . . . *axiom.*

hyp. . . . . *hypotheſis.* It may be neceſſary here to remark, that the *hypotheſis* is the condition aſſumed or taken for granted. Thus, the hypotheſis of the propoſition given in the Introduction, is that the triangle is iſoſceles, or that its legs are equal.

conſt. . . . . *conſtruction.* The *conſtruction* is the change made in the original figure, by drawing lines, making angles, deſcribing circles, &c. in order to adapt it to the argument of the demonſtration or the ſolution of the problem. The conditions under which theſe changes are made, are as indisputable as thoſe contained in the hypotheſis. For inſtance, if we make an angle equal to a given angle, theſe two angles are equal by conſtruction.

Q. E. D. . . . . *Quod erat demonſtrandum.*
Which was to be demonſtrated.

Harvard University - Houghton Library / pga_typ_805_47_3730

Faults to be corrected before reading this Volume.

PAGE 13, line 9, *for* def. 7 *read* def. 10.

45, laſt line, *for* pr. 19 *read* pr. 29.

54, line 4 from the bottom, *for* black and red line *read* blue and red line.

59, line 4, *for* add black line ſquared *read* add blue line ſquared.

60, line 17, *for* red line multiplied by red and yellow line *read* red line multiplied by red, blue, and yellow line.

76, line 11, *for* def. 7 *read* def. 10.

81, line 10, *for* take black line *read* take blue line.

105, line 11, *for* yellow black angle add blue angle equal red angle *read* yellow black angle add blue angle add red angle.

129, laſt line, *for* circle *read* triangle.

141, line 1, *for* Draw black line *read* Draw blue line.

196, line 3, before the yellow magnitude inſert M.


Harvard University - Houghton Library / pga_typ_805_47_3730

Harvard University - Houghton Library / pga_typ_805_47_3730

Euclid.

BOOK I.

PROPOSITION I. PROBLEM.



ON *a given finite ſtraight line* (——) *to deſcribe an equilateral triangle.*

Deſcribe ◯ and ◯ (poſtulate 3.); draw —— and —— (poſt. 1.).

then will △ be equilateral.

For —— = —— (def. 15.);


and —— = —— (def. 15.),

∴ —— = —— (axiom. 1.);

and therefore △ is the equilateral triangle required.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



FROM *a given point* (—), *to draw a ſtraight line equal to a given finite ſtraight line* (———).


Draw ---------- (poſt. 1.), deſcribe △ (pr. 1.), produce —— (poſt. 2.), deſcribe ◯ (poſt. 3.), and ◯ (poſt. 3.); produce —— (poſt. 2.), then —— is the line required.

For ——— = ——— (def. 15.), and —— = —— (conſt.), ∴ ——— = ——— (ax. 3.), but (def. 15.) ——— = ——— = ———; ∴ ——— drawn from the given point (——), is equal the given line ———.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

FROM *the greater* (——·····) *of two given ſtraight lines, to cut off a part equal to the leſs* (——).



Draw —— = —— (pr. 2.); deſcribe



(poſt. 3 .), then —— = ——.


For —— = —— (def. 15.),

and —— = —— (conſt.);

∴ —— = —— (ax. 1.).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730




IF *two triangles have two ſides of the one reſpectively equal to two ſides of the other,* (—— *to* —— *and* —— *to* ——) *and the angles* (▲ *and* ▲) *contained by thoſe equal ſides alſo equal; then their baſes or their ſides* (—— *and* ——) *are alſo equal: and the remaining and their remaining angles oppoſite to equal ſides are reſpectively equal* (◣ = ◣ *and* ▲ = ▲): *and the triangles are equal in every reſpect.*

Let the two triangles be conceived, to be ſo placed, that the vertex of the one of the equal angles, ▲ or ▲ ; ſhall fall upon that of the other, and —— to coincide with ——, then will —— coincide with —— if applied: conſequently —— will coincide with ——, or two ſtraight lines will encloſe a ſpace, which is impoſſible (ax. 10), therefore —— = ——, ◣ = ◣ and ▲ = ▲, and as the triangles  and  coincide, when applied, they are equal in every reſpect.


Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730


IN *any iſoſceles triangle* △ *if the equal ſides be produced, the external angles at the baſe are equal, and the internal angles at the baſe are alſo equal.*






Produce ——, and ——, (poſt. 2.), take —— = ——, (pr. 3.); draw —— and ——.

Then in  and  we have, —— = —— (conſt.),  common to both, and —— = —— (hyp.) ∴  , —— = —— and  (pr. 4.).

Again in  and we have —— = —— ,

 and —— = —— , ∴

 and  (pr. 4.) but

 , ∴  (ax. 3.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IN *any triangle* (△) *if two angles* (▲ *and* ◢) *are equal, the ſides* (▬·▬· *and* ▬▬) *oppoſite to them are alſo equal.*


For if the ſides be not equal, let one of them ▬·▬· be greater than the other ▬▬, and from it cut off ▬▬ = ▬▬ (pr. 3.), draw ▬▬.

Then in △ and △, ▬▬ = ▬▬, (conſt.) ▲ = ◢ (hyp.) and ▬▬ common, ∴ the triangles are equal (pr. 4.) a part equal to the whole, which is abſurd; ∴ neither of the ſides ▬·▬· or ▬▬ is greater than the other, ∴ hence they are equal

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

ON *the ſame baſe* (———), *and on the ſame ſide of it there cannot be two triangles having their conterminous ſides* (——— *and* ———, ——— *and* ———) *at both extremities of the baſe, equal to each other.*




When two triangles ſtand on the ſame baſe, and on the ſame ſide of it, the vertex of the one ſhall either fall outſide of the other triangle, or within it; or, laſtly, on one of its ſides.

If it be poſſible let the two triangles be conſtructed ſo that { ——— = ——— ; ——— = ——— }, then draw ------ and,




$\therefore$ 

but (pr. 5.)  } which is abſurd,

therefore the two triangles cannot have their conterminous ſides equal at both extremities of the baſe.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *two triangles have two ſides of the one reſpectively equal to two ſides of the other* (——— = ——— *and* ——— = ———), *and alſo their baſes* (——— = ———), *equal; then the angles* (◀ *and* ◀) *contained by their equal ſides are alſo equal.*

If the equal baſes ——— and ——— be conceived to be placed one upon the other, ſo that the triangles ſhall lie at the ſame ſide of them, and that the equal ſides ——— and ———, ——— and ——— be conterminous, the vertex of the one muſt fall on the vertex of the other; for to ſuppoſe them not coincident would contradict the laſt propoſition.


Therefore the ſides ——— and ———, being coincident with ——— and ———,

$\therefore$ ▲ = ▲.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *biſect a given rectilinear angle* ( ).



Take —— = —— (pr. 3.)

draw ——, upon which

deſcribe ▽ (pr. 1.),

draw ——.

Becauſe —— = —— (conſt.)

and —— common to the two triangles

and —— = —— (conſt.),

∴ ◢ = ◣ (pr. 8.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *biſect a given finite ſtraight line* (——·····).

Conſtruct △ (pr. 1.),

draw ——, making ◔ = ◔ (pr. 9.).

Then —— = ····· by (pr. 4.),

for —— = —— (conſt.) ◔ = ◔


and —— common to the two triangles.

Therefore the given line is biſected.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

FROM *a given point* (—·—), *in a given ſtraight line* (———), *to draw a perpendicular.*



Take any point (— — —) in the given line,

cut off ——— = ——— (pr. 3.),


conſtruct  (pr. 1.),

draw ——— and it ſhall be perpendicular to the given line.

For ——— = ——— (conſt.)

——— = ——— (conſt.)


and ——— common to the two triangles.

Therefore  (pr. 8.)

∴ ——— ⊥ ——— (def. 10.).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *draw a ſtraight line perpendicular to a given indefinite ſtraight line* (—) *from a given* (*point* ∧) *without.*

With the given point ∧ as centre, at one ſide of the line, and any diſtance — capable of extending to the other ſide, deſcribe ⌒,

Make — = — (pr. 10.)

draw —, — and —.

then — ⊥ —.

For (pr. 8.) ſince — = — (conſt.)

— common to both,


and — = — (def. 15.)

∴ ◔ = ◔, and

∴ — ⊥ — (def. 10.).


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



WHEN *a ſtraight line* (——) *ſtanding upon another ſtraight line* (——) *makes angles with it; they are either two right angles or together equal to two right angles.*


If —— be ⊥ to —— then,



(def. 7.),

But if —— be not ⊥ to ——,

draw —— ⊥ ——; (pr. 11.)



(conſt.),



∴ 
(ax. 2.)



Q. E. D.



IF *two ſtraight lines* (—— *and* ——), *meeting a third ſtraight line* (——), *at the ſame point, and at oppoſite ſides of it, make with it adjacent angles* (*and*) *equal to two right angles; theſe ſtraight lines lie in one continuous ſtraight line.*

For, if poſſible let ——, and not ——,
be the continuation of ——,

then + = 

but by the hypotheſis + =

∴ = , (ax. 3.); which is abſurd (ax. 9.).

∴ ——, is not the continuation of ——, and the like may be demonſtrated of any other ſtraight line except ——, ∴ —— is the continuation of ——.


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *two right lines* (—— *and* ——) *interſect one another, the vertical angles*  *and*  , 


and  *are equal.*






∴ .

In the ſame manner it may be ſhown that



Q. E. D.



IF *a ſide of a trian-*  *gle* ( ) *is produced, the external angle* ( ) *is greater than either of the internal remote angles* (*or*).

Make —— = - - - - - - (pr. 10.).

Draw —— and produce it until

- - - - - - = ——; draw ——.

In  and  ; —— = - - - - - -

 =  and —— = - - - - - -

(conſt. pr. 15.), ∴  =  (pr. 4.),

 ⊏  .

In like manner it can be ſhown, that if - - - - - - be produced,  ⊏  , and therefore  which is =  is ⊏  .

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730

ANY *two angles of a tri-*




angle are to-


gether leſs than two right angles.



Produce ——, then will




But  (pr. 16.)

$\therefore$  ,

and in the ſame manner it may be ſhown that any other two angles of the triangle taken together are leſs than two right angles.

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730





IN *any triangle*  *if one ſide* ····—— *be greater than another* ——, *the angle oppoſite to the greater ſide is greater than the angle oppoſite to the leſs.*

i. e. 

Make —— = —— (pr. 3.), draw ——.

Then will  (pr. 5.);

but  (pr. 16.)

∴ 

and much more

is  .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *in any triangle* △ *one angle* ◗ *be greater than another* ◢ *the ſide* ——— *which is oppoſite to the greater angle, is greater than the ſide* —— *oppoſite the leſs.*



If ——— be not greater than —— then muſt ——— = or ⊐ ——.

If ——— = —— then ◗ = ◢ (pr. 5.);

which is contrary to the hypotheſis.

——— is not leſs than ——; for if it were,


◗ ⊐ ◢ (pr. 18.)

which is contrary to the hypotheſis:

∴ ——— ⊏ ——.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



ANY *two ſides* —— *and* —— *of a triangle* △ *taken together are greater than the third ſide* (——).

Produce ——, and

make - - - - - = —— (pr. 3.);

draw ——.

Then becauſe - - - - - = —— (conſt.),

◔ = ◔ (pr. 5.)

∴ ◔ ⊏ ◔ (ax. 9.)

∴ —— + - - - - - ⊏ —— (pr. 19.)


and ∴ —— + —— ⊏ ——.

Q. E. D

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *from any point* ( ) *within a triangle* *ſtraight lines be drawn to the extremities of one ſide* (_____), *theſe lines muſt be together leſs than the other two ſides, but muſt contain a greater angle.*





Produce ——,

—— + —— ⊏ ——····· (pr. 20.),

add ----- to each,

—— + ——····· ⊏ ——···· + ------ (ax. 4.)

In the ſame manner it may be ſhown that

——···· + ----- ⊏ —— + ——, ∴

—— + ——···· ⊏ —— + ——,

which was to be proved.


Again  (pr. 16.),

and alſo  (pr. 16.),

∴  .

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730



GIVEN *three right lines*  *the ſum of any two greater than the third, to conſtruct a triangle whoſe ſides ſhall be reſpectively equal to the given lines.*

Aſſume  (pr. 3.).


Draw 
and  } (pr. 2.).

With  and as radii,

deſcribe  and  (poſt. 3.);

draw  and ,


then will  be the triangle required.

For  (conſt.)


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

AT *a given point* ( ) *in a given ſtraight line* ( ), *to make an angle equal to a given rectilineal angle* ( ).



Draw —— between any two points in the legs of the given angle.



Conſtruct  (pr. 22.).


ſo that —— = ——, —— = ——

and —— ——.

Then  =  (pr. 8.).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730





IF *two triangles have two ſides of the one reſpectively equal to two ſides of the other* (—— *to* —— *and* - - - - - - *to* ——), *and if one of the angles* () *contained by the equal ſides be greater than the other* (), *the ſide* (——) *which is oppoſite to the greater angle is greater than the ſide* (——) *which is oppoſite to the leſs angle.*

Make  =  (pr. 23.),

and —— = —— (pr. 3.),

draw ..... and - - - - - - - -.

Becauſe —— = - - - - - - - - (ax. 1. hyp. conſt.)

∴  =  (pr. 5.)

but  ⊐ 

and ∴  ⊐  ,


∴ —— ⊏ - - - - - - - - (pr. 19.)

but - - - - - - - - = —— (pr. 4.)

∴ —— ⊏ ——.

Q. E. D.

IF *two triangles have two ſides* (——— *and* ———) *of the one reſpectively equal to two ſides* (——— *and* ———) *of the other, but their baſes unequal, the angle ſubtended by the greater baſe* (———) *of the one, muſt be greater than the angle ſubtended by the leſs baſe* (———) *of the other.*



△ =, ⊏ or ⊐ △ △ is not equal to △

for if △ = △ then ——— = ——— (pr. 4.)

which is contrary to the hypotheſis;

△ is not leſs than △

for if △ ⊐ △

then ——— ⊐ ——— (pr. 24.),

which is alſo contrary to the hypotheſis:

∴ △ ⊏ △.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

CASE I.



CASE II.



IF *two triangles have two angles of the one respectively equal to two angles of the other,* (= *and* =), *and a side of the one equal to a side of the other similarly placed with respect to the equal angles, the remaining sides and angles are respectively equal to one another.*

CASE I.

Let —— and —— which lie between the equal angles be equal,

then —— = ——.

For if it be possible, let one of them —— be greater than the other;


make —— = ——, draw ——.


In and we have

— = —, = , — = —;


∴ = (pr. 4.)

but  (hyp.)


and therefore  , which is abſurd;
hence neither of the ſides and is greater than the other; and ∴ they are equal;


∴ , and  , (pr. 4.).


CASE II.


Again, let  , which lie oppoſite the equal angles  and  . If it be poſſible, let  , then take  , draw  .

Then in  and  we have  ,

 and  ,

∴  (pr. 4.)

but  (hyp.)

∴  which is abſurd (pr. 16.).

Conſequently, neither of the ſides  or  is greater than the other, hence they muſt be equal. It follows (by pr. 4.) that the triangles are equal in all reſpects.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730




IF *a ſtraight line* (——) *meeting two other ſtraight lines,* (—— *and* ——) *makes with them the alternate angles* (and ; and) *equal, theſe two ſtraight lines are parallel.*

If —— be not parallel to —— they ſhall meet when produced.

If it be poſſible, let thoſe lines be not parallel, but meet when produced; then the external angle is greater than (pr. 16), but they are alſo equal (hyp.), which is abſurd: in the ſame manner it may be ſhown that they cannot meet on the other ſide; ∴ they are parallel.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *a ſtraight line* (——), *cutting two other ſtraight lines* (—— *and* ——), *makes the external equal to the internal and oppoſite angle, at the ſame ſide of the cutting line* (*namely,* = *or* =), *or if it makes the two internal angles at the ſame ſide* (*and* , *or* *and*) *together equal to two right angles, thoſe two ſtraight lines are parallel.*

Firſt, if = , then = (pr. 15.),

∴ = ∴ —— || —— (pr. 27.).

Secondly, if + = ,

then + = (pr. 13.),


∴ + = + (ax. 3.)

∴ =

∴ —— || —— (pr. 27.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



A STRAIGHT *line* (——) *falling on two parallel ſtraight lines* (—— *and* ——), *makes the alternate angles equal to one another; and alſo the external equal to the internal and oppoſite angle on the ſame ſide; and the two internal angles on the ſame ſide together equal to two right angles.*

For if the alternate angles  and  be not equal,

draw ——, making  =  (pr. 23).

Therefore —— ‖ —— (pr. 27.) and therefore two ſtraight lines which interſect are parallel to the ſame ſtraight line, which is impoſſible (ax. 12).

Hence the alternate angles  and  are not unequal, that is, they are equal:  =  (pr. 15);

∴  =  , the external angle equal to the internal and oppoſite on the ſame ſide: if  be added to

both, then 


(pr. 13).

That is to ſay, the two internal angles at the ſame ſide of the cutting line are equal to two right angles.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

STRAIGHT *lines* ( ) *which are parallel to the same ſtraight line* ( ), *are parallel to one another.*



Let ——— interſect { } ;

Then, ▲ = ▲ = ▲ (pr. 29.),

∴ ▲ = ▲

∴ ——— ‖ ——— (pr. 27.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



FROM *a given point* 7 *to draw a ſtraight line parallel to a given ſtraight line* (——).

Draw —— from the point 7 to any point ∠ in ——,


make ▽ = ◠ (pr. 23.),

then —·— ∥ —— (pr. 27.).


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730


IF *any side* (——) *of a triangle be produced, the external angle* () *is equal to the sum of the two internal and opposite angles* ( *and* ), *and the three internal angles of every triangle taken together are equal to two right angles.*




Through the point  draw

 (pr. 31.).

Then  (pr. 29.),

$\therefore$  (ax. 2.),


and therefore



(pr. 13.).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



STRAIGHT *lines* (——— *and* ———) *which join the adjacent extremities of two equal and parallel ſtraight lines* (——— *and* ----------), *are themſelves equal and parallel.*

Draw ——— the diagonal.

——— = ---------- (hyp.)

▼ = ▲ (pr. 29.)

and ——— common to the two triangles;

∴ ——— = ———, and ▼ = ▲ (pr. 4.);

and ∴ ——— ‖ ——— (pr. 27.).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

THE *oppoſite ſides and angles of any parallelogram are equal, and the diagonal* (——) *divides it into two equal parts.*



Since { = , = } (pr. 29.)

and —— common to the two triangles.

∴ { = , = , = } (pr. 26.)

and = (ax.):

Therefore the oppoſite ſides and angles of the parallelogram are equal: and as the triangles  and  are equal in every reſpect (pr. 4,), the diagonal divides the parallelogram into two equal parts.


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



PARALLELOGRAMS *on the ſame baſe, and between the ſame parallels, are (in area) equal.*

On account of the parallels,










Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

PARALLELO-
GRAMS
( *and* ) *on equal baſes, and between the ſame parallels, are equal.*



Draw —— and - - - - - -,

—— = —— = ——, by (pr. 34, and hyp.);

∴ —— = and || ——;

∴ —— = and || - - - - - - - (pr. 33.)


And therefore  is a parallelogram:


but  (pr. 35.)

∴  (ax. 1.).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TRIANGLES  *and* *on the ſame baſe* (——) *and between the ſame parallels are equal.*

Draw ------ ‖ ——
····· ‖ ——
} (pr. 31.)

Produce ----------.

 and  are parallelograms on the ſame baſe, and between the ſame parallels, and therefore equal. (pr. 35.)

∴ {  = twice 
 = twice  } (pr. 34.)


∴  =  .


Q. E D.


Harvard University - Houghton Library / pga_typ_805_47_3730


TRIANGLES  ( *and* ) *on equal baſes and between the ſame parallels are equal.*

Draw  (pr. 31.).

 (pr. 36.);


but  (pr. 34.),

and  (pr. 34.),

 (ax. 7.).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730




EQUAL triangles  *and*  *on the ſame baſe* (——) *and on the ſame ſide of it, are between the ſame parallels.*

If ——, which joins the vertices of the triangles, be not ‖ ——, draw —— ‖ —— (pr. 31.), meeting - - - - - - -.

Draw ——.

Becauſe —— ‖ —— (conſt.)

 (pr. 37.):

but  (hyp.);


 , a part equal to the whole, which is abſurd.

∴ —— ≠ ——; and in the ſame manner it can be demonſtrated, that no other line except —— is ‖ ——; ∴ —— ‖ ——.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

EQUAL *triangles* (△ *and* △) *on equal baſes, and on the ſame ſide, are between the ſame parallels.*



If —— which joins the vertices of triangles be not ‖ —— ,
draw —— ‖ —— (pr. 31.),
meeting ---------.

Draw ——.

Becauſe —— ‖ —— (conſt.)

△ = △ but △ = △

∴ △ = △, a part equal to the whole, which is abſurd.

∴ —— ∦ ——: and in the ſame manner it can be demonſtrated, that no other line except —— is ‖ ——: ∴ —— ‖ ——.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *a parallelogram*  *and a triangle*  *are upon the ſame baſe* ——— *and between the ſame parallels* - - - - - - *and* ———, *the parallelogram is double the triangle.*

Draw —— the diagonal;

Then  =  (pr. 37.)


 = twice  (pr. 34.)

∴  = twice  .


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730


TO *conſtruct a parallelogram equal to a given*




triangle  *and having an angle equal to a given rectilinear angle*  .

Make  (pr. 10.)


Draw  .

Make  (pr. 23.)

Draw  (pr. 31.)


 (pr. 41.)

but  (pr. 38.)

∴  .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



THE *complements*  *and*  *of the parallelograms which are about the diagonal of a parallelogram are equal.*

 (pr. 34.)

and  (pr. 34.)

$\therefore$  (ax. 3.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *a given ſtraight line* (——) *to apply a parallelogram equal to a given triangle* ( ), *and having an angle equal to a given rectilinear angle* ( ).



Make = (pr. 42.) with =

and having one of its ſides ------ conterminous with and in continuation of ——.

Produce —— till it meets —— || ------

draw —— produce it till it meets ------ continued;

draw ------ || ·····—— meeting ——

produced, and produce ·····.

= (pr. 43.)

but = (conſt.)

∴ = ; and

= = = (pr. 19. and conſt.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *construct a parallelogram equal to a given rectilinear figure* ( ) *and having an angle equal to a given rectilinear angle* ( ).

Draw —— and —— dividing the rectilinear figure into triangles.



Construct =

having = (pr. 42.)

to —— apply =

having = (pr. 44.)

to —— apply =

having = (pr. 44.)

∴ =

and is a parallelogram. (prs. 29, 14, 30.)

having = .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

UPON *a given ſtraight line* (——) *to conſtruct a ſquare.*



Draw —— ⊥ and = ——
(pr. 11. and 3.)

Draw —— || ——, and meeting —— drawn || ——.

In  —— = —— (conſt.)


 = a right angle (conſt.)


∴  =  = a right angle (pr. 29.),
and the remaining ſides and angles muſt be equal, (pr. 34.)

and ∴  is a ſquare. (def. 27.)

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730



IN *a right angled triangle*  *the ſquare on the hypotenuſe* —— *is equal to the ſum of the ſquares of the ſides,* (—— *and* ——).


On ——, —— and ——
deſcribe ſquares, (pr. 46.)

Draw ---------- ‖ - - - - - - (pr. 31.)
alſo draw —— and ——.


 ,

To each add  ∴  ,

—— = - - - - - - and —— = ----------;


∴  .


Again, becauſe —— ‖ ----------





In the ſame manner it may be ſhown





Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *the ſquare of one ſide* (——) *of a triangle is equal to the ſquares of the other two ſides* (—— *and* ——), *the angle* ( ) *ſubtended by that ſide is a right angle.*

Draw ---------- ⊥ —— and = —— (prs. 11. 3.)
and draw ---------- alſo.

Since ---------- = —— (conſt.)


$\therefore$ ------^{2} + ——^{2} = ——^{2} + ——^{2},

but ------^{2} + ——^{2} = ------^{2} (pr. 47.),

and ——^{2} + ——^{2} = ——^{2} (hyp.)

$\therefore$ ------^{2} = ——^{2},

$\therefore$ ------ = ——;

and $\therefore$  = (pr. 8.),

consequently  is a right angle.


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

BOOK II.

DEFINITION I.

A RECTANGLE or a right angled parallelogram is ſaid to be contained by any two of its adjacent or conterminous ſides.



Thus: the right angled parallelogram  is ſaid to be contained by the ſides ——— and ———;

or it may be briefly deſignated by

——— · ———.


If the adjacent ſides are equal; i. e. ——— = ———,

then ——— · ——— which is the expreſſion for the rectangle under ——— and ———

is a ſquare, and

is equal to { ——— · ——— or ———²
{ ——— · ——— or ———²


DEFINITION II.




IN a parallelogram, the figure compoſed of one of the parallelograms about the diagonal, together with the two complements, is called a *Gnomon*.

Thus  and  are called Gnomons.

Harvard University - Houghton Library / pga_typ_805_47_3730



THE *rectangle contained by two ſtraight lines, one of which is divided into any number of parts,*



is equal to the ſum of the rectangles contained by the undivided line, and the ſeveral parts of the divided line.


Draw  and  (prs. 2. 3. B. 1.); complete the parallelograms, that is to ſay,

Draw  (pr. 31. B. 1.)











Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730




IF *a ſtraight line be divided into any two parts* ——, *the ſquare of the whole line is equal to the ſum of the rectangles contained by the whole line and each of its parts.*







Deſcribe (B. 1. pr. 46.)

Draw —— parallel to - - - - - - (B. 1. pr. 31.)












Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *a ſtraight line be divided into any two parts* —— —, *the rectangle contained by the whole line and either of its parts, is equal to the ſquare of that part, together with the rectangle under the parts.*



—— — . —— = —— ^{2} + —— . —, or,

—— —. —— = — ^{2} + —— .—.

Deſcribe  (pr. 46, B. 1.)

Complete  (pr. 31, B. 1.)

Then  =  +  , but

 = —— . —— and


 = —— ^{2},  = —— . —,

∴ —— —. —— = —— ^{2} + —— . —:

In a ſimilar manner it may be readily ſhown that —— —. — = — ^{2} + —— . —.

Q. E. D

Harvard University - Houghton Library / pga_typ_805_47_3730




IF *a ſtraight line be divided into any two parts* —— —, *the ſquare of the whole line is equal to the ſquares of the parts, together with twice the rectangle contained by the parts.*

—— —^{2} = ——^{2} + —^{2} +
twice —— · —.

Deſcribe  (pr. 46, B. 1.)

draw —————— (poſt. 1.),

and { —·· || ——
      ——— || ——· } (pr. 31, B. 1.)
∴

 (pr. 5, B. 1.),

 (pr. 29, B. 1.)

∴ 


Harvard University - Houghton Library / pga_typ_805_47_3730


∴ by (prs. 6, 29, 34. B. 1.)  is a ſquare = .

For the ſame reaſons  is a ſquare =  ,

 =  =  (pr. 43, b. 1.)


but  =  +  +  +  ,

∴  = +  +


twice  .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730




IF *a ſtraight line be divided into two equal parts and alſo into two unequal parts, the rectangle contained by the unequal parts, together with the ſquare of the line between the points of ſection, is equal to the ſquare of half that line*



Deſcribe  (pr. 46, B. 1.), draw and



(pr. 31, B. 1.)


(p. 36, B. 1.)


(p. 43, B. 1.)


∴ (ax. 2.) 

Harvard University - Houghton Library / pga_typ_805_47_3730

but  =  (cor. pr. 4. B. 2.)

and  = (conft.)


∴ (ax. 2.)  = 

∴ 

.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730




IF *a ſtraight line be biſected* — — *and produced to any point* — — —, *the rectangle contained by the whole line ſo increaſed, and the part produced, together with the ſquare of half the line, is equal to the ſquare of the line made up of the half, and the produced part.*

— — · — + — —^{2} = — — —^{2}.

Deſcribe  (pr. 46, B. 1.), draw —————

and { — - - - = — - - - ; - - - - - - - - - - — = — — — — — ; - - - - = — - - - } (pr. 31, B. 1.)

 (prs. 36, 43, B. 1)

∴  = — · — —;

but  = —^{2} (cor. 4, B. 2.)

∴  = — —^{2} =  (conſt. ax. 2.)

∴ — — — · — + —^{2} = — — —^{2}.


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *a ſtraight line be divided into any two parts* —— —, *the ſquares of the whole line and one of the parts are equal to twice the rectangle contained by the whole line and that part, together with the ſquare of the other parts.*

—— —^{2} + —^{2} =


2 —— — · — + ——^{2}




Deſcribe  , (pr. 46, B. 1.).

Draw ——————— (poſt. 1.),

and { —···· ‖ ——— ; ·——— ‖ —— — } (pr. 31, B. 1.).

 (pr. 43, B. 1.),

add  = —^{2} to both, (cor. 4, B. 2.)



 = ——^{2} (cor. 4, B. 2.)


∴ 

——^{2} =  ;

∴ ——— —^{2} + —^{2} = 2 ——— · — + ——^{2}.

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730



IF *a ſtraight line be divided into any two parts* —, *the ſquare of the ſum of the whole line and any one of its parts, is equal to four times the rectangle contained by the whole line, and that part together with the ſquare of the other part.*

—^{2} = 4 · — · — + —^{2},

Produce — and make — = —

Conſtruct  (pr. 46, B. 1.);

draw ——,

— } ∥ — } (pr. 31, B. 1.)

— } ∥ — }

—^{2} = —^{2} + —^{2} + 2 · — · —

(pr. 4, B. 11.)

but —^{2} + —^{2} = 2 · — · — + —^{2}


(pr. 7, B. 11.)

∴ —^{2} = 4 · — · — + —^{2}.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *a ſtraight line be divided into two equal parts* ——, *and alſo into two unequal parts* ——, *the ſquares of the unequal parts are together double the ſquares of half the line, and of the part between the points of ſection.*





—— ^{2} + —— ^{2} = 2 —— ^{2} + 2 —— ^{2}.

Make —— ⊥ and = —— or ——,


Draw ---------- and ----------,

—— ‖ ——, —— ‖ ——, and draw ——.

 = (pr. 5, B. 1.) = half a right angle.
(cor. pr. 32, B. 1.)

 = (pr. 5, B. 1.) = half a right angle.
(cor. pr. 32, B. 1.)

∴  = a right angle.



(prs. 5, 29, B. 1.).

hence —— = ——, ---- = —— = ——


(prs. 6, 34, B. 1.)

$$——^2 = \begin{cases} ——^2 + ——^2, \text{ or } + ——^2 \\ = \begin{cases} ----------^2 = 2\ ——^2 \\ \text{(pr. 47, B. 1.)} \\ ----------^2 = 2\ ——^2 \end{cases} \end{cases}$$

∴ ——^{2} + ——^{2} = 2 ——^{2} + 2 ——^{2}.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *a ſtraight line* —— *be biſected and produced to any point* ——, *the ſquares of the whole produced line, and of the produced part, are together double of the ſquares of the half line, and of the line made up of the half and produced part.*


—— ^{2} + —— ^{2} = 2 —— ^{2} + 2 —— ^{2}.

Make ═ ⊥ and = to —— or ——,

draw ——···· and ——----,

and { —— ‖ ═ ; ——— ‖ —— } (pr. 31, B. 1.);

draw —— alſo.

 = (pr. 5, B. 1.) = half a right angle. (cor. pr. 32, B. 1.)


 =  (pr. 5, B. 1.) = half a right angle (cor. pr. 32, B. 1.)

∴  = a right angle.


Harvard University - Houghton Library / pga_typ_805_47_3730




half a right angle (prs. 5, 32, 29, 34, B. 1.),

and 

(prs. 6, 34, B. 1.). Hence by (pr. 47, B. 1.)



Q. E. D.



TO *divide a given ſtraight line* ▬▬ *in ſuch a manner, that the rectangle contained by the whole line and one of its parts may be equal to the ſquare of the other.*

▬▬ · ▬▬ = ▬▬^{2}.

Deſcribe  (pr. 46, B. 1.),

make ▬ = ▬▬ (pr. 10, B. 1.),

draw ▬▬,


take ▬ = ▬▬ (pr. 3, B. 1.),

on ▬ deſcribe  (pr. 46, B. 1.),

Produce ▬▬▬▬▬▬ (poſt. 2.).

Then, (pr. 6, B. 2.) ▬ · ▬ + ▬^{2} = ▬^{2} = ▬▬^{2} = ▬^{2} +

▬^{2} ∴ ▬ · ▬ = ▬^{2}, or,




▬ · ▬ = ▬^{2}.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IN *any obtuſe angled triangle, the ſquare of the ſide ſubtending the obtuſe angle exceeds the ſum of the ſquares of the ſides containing the obtuſe angle, by twice the rectangle contained by either of theſe ſides and the produced parts of the ſame from the obtuſe angle to the perpendicular let fall on it from the oppoſite acute angle.*



——² ⊏ ——² + ——² by

2 —— · ------.

By pr. 4, B. 2.

——.....² = ——² + ---------² + 2 —— · -----:

add ——² to both

------² + ——² = ——² (pr. 47, B. 1.)

= 2 · —— · ------- + ——² + { ---------² / ——² } or

+ ——² (pr. 47, B. 1.). Therefore,

——² = 2 · —— · --------- + ——² +

——²: hence ——² ⊏ ——² + ——²

by 2 · —— · ------.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730


FIRST.

SECOND.




IN *any triangle, the ſquare of the ſide ſubtending an acute angle, is leſs than the ſum of the ſquares of the ſides containing that angle, by twice the rectangle contained by either of theſe ſides, and the part of it intercepted between the foot of the perpendicular let fall on it from the oppoſite angle, and the angular point of the acute angle.*


FIRST.




SECOND.




Firſt, ſuppoſe the perpendicular to fall within the triangle, then (pr. 7, B. 2.)



add to each  then,



∴ (pr. 47, B. 1.)



and $\therefore$ $\overline{AC}^2 \sqsupset \overline{BC}^2 + \overline{AB}^2$ by $2 \cdot \overline{BC} \cdot \overline{BD}$.

Next ſuppoſe the perpendicular to fall without the triangle, then (pr. 7, B. 2.)

$$\overline{AC}^2 + \overline{BC}^2 = 2 \cdot \overline{AC} \cdot \overline{BC} + \overline{AB}^2,$$

add to each $\overline{BC}^2$ then


$$\overline{AC}^2 + \overline{BC}^2 + \overline{BC}^2 = 2 \cdot \overline{AC} \cdot \overline{BC} + \overline{AB}^2 + \overline{BC}^2 \therefore \text{(pr. 47, B. 1.)},$$

$$\overline{AB}^2 + \overline{BC}^2 = 2 \cdot \overline{AC} \cdot \overline{BC} + \overline{AB}^2,$$

$\therefore \overline{AC}^2 \sqsupset \overline{AB}^2 + \overline{BC}^2$ by $2 \cdot \overline{AC} \cdot \overline{BC}$.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *draw a right line of which the ſquare ſhall be equal to a given rectilinear figure.*

To draw ——— *ſuch that,*

$$\text{———}^2 = \text{[figure]}$$

Make [figure] = [figure] (pr. 45, B. 1.),

produce ····——— until ····· = — ;

take ·····—— = ——— (pr. 10, B. 1.),

Deſcribe [figure] (poſt. 3.),

and produce — to meet it: draw ———.

$$\text{———}^2 \text{ or } \text{———}^2 = \text{·····}\cdot\text{·····——} + \text{·····}^2$$

(pr. 5, B. 2.),

but $\text{———}^2 = \text{———}^2 + \text{·····}^2$ (pr. 47, B. 1.);

$\therefore \text{———}^2 + \text{·····}^2 = \text{·····}\cdot\text{·····——} + \text{·····}^2$,

$\therefore \text{———}^2 = \text{·····}\cdot\text{·····——}$, and

$\therefore \text{———}^2 =$ [figure] $=$ [figure]

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

BOOK III.


DEFINITIONS.

I.

EQUAL circles are thoſe whoſe diameters are equal.


II.

A right line is said to touch a circle when it meets the circle, and being produced does not cut it.




III.

Circles are ſaid to touch one another which meet but do not cut one another.



IV.

Right lines are ſaid to be equally diſtant from the centre of a circle when the perpendiculars drawn to them from the centre are equal.



Harvard University - Houghton Library / pga_typ_805_47_3730

V.


And the ſtraight line on which the greater perpendicular falls is ſaid to be farther from the centre.



VI.


A ſegment of a circle is the figure contained by a ſtraight line and the part of the circumference it cuts off.

VII.



An angle in a ſegment is the angle contained by two ſtraight lines drawn from any point in the circumference of the ſegment to the extremities of the ſtraight line which is the baſe of the ſegment.

VIII.



An angle is ſaid to ſtand on the part of the circumference, or the arch, intercepted between the right lines that contain the angle.

IX.



A ſector of a circle is the figure contained by two radii and the arch between them.

Harvard University - Houghton Library / pga_typ_805_47_3730

X.


Similar ſegments of circles are thoſe which contain equal angles.





Circles which have the ſame centre are called *concentric circles*.






TO *find the centre of a given circle* ◯.


Draw within the circle any ſtraight line ——·····, make —— = ·····, draw —— ⊥ ——·—·—; biſect ——, and the point of biſection is the centre.

For, if it be poſſible, let any other point as the point of concourſe of ——, ····· and ▪▪▪▪▪▪▪▪▪ be the centre.

Becauſe in  and 

—— = ▪▪▪▪▪▪ (hyp. and B. 1, def. 15.)


—— = ·—·—· (conſt.) and ····· common,

 =  (B. 1, pr. 8.), and are therefore right angles; but  =  (conſt.)  =  (ax. 11.)

which is abſurd; therefore the aſſumed point is not the centre of the circle; and in the ſame manner it can be proved that no other point which is not on —— is the centre, therefore the centre is in ——, and therefore the point where —— is biſected is the centre.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



A STRAIGHT *line* (——) *joining two points in the circumference of a circle*



, *lies wholly within the circle.*

Find the centre of  (B.3.pr.1.);

from the centre draw —— to any point in ——,

meeting the circumference from the centre;

draw —— and ——.

Then ◥ = ◥ (B. 1. pr. 5.)

but ▲ ⊏ ◥ or ⊏ ◥ (B. 1. pr. 16.)

∴ —— ⊏ —— (B. 1. pr. 19.)

but —— = ——-----,


∴ ——----- ⊏ ——;

∴ —— ⊐ ——----- ;

∴ every point in —— lies within the circle.

Q. E. D

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *a ſtraight line* (——) *drawn through the centre of a circle* (○) *biſects a chord* (——····) *which does not paſs through the centre, it is perpendicular to it; or, if perpendicular to it, it biſects it.*

Draw —— and —— to the centre of the circle.

In △ and △,

—— = ——, —— common, and

—— = ····· ∴ ◔ = ◔ (B. 1. pr. 8.)

and ∴ —— ⊥ ——···· (B. 1. def. 7.)

Again let —— ⊥ ——··

Then in △ and △

▲ = ▲ (B. 1. pr. 5.)

◔ = ◔ (hyp.)

and —— = ——


∴ —— = ····· (B. 1. pr. 26.)

and ∴ —— biſects ——···.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730


IF *in a circle two ſtraight lines cut one another, which do not both paſs through the centre, they do not biſect one another.*




If one of the lines paſs through the centre, it is evident that it cannot be biſected by the other, which does not paſs through the centre.

But if neither of the lines ——— or ——— paſs through the centre, draw - - - - - - - from the centre to their interſection.

If ——— be biſected, - - - - - - ⊥ to it (B. 3. pr. 3.)

∴  and if ——— be

biſected, - - - - - - ⊥ ——— (B. 3. pr. 3.)

∴ 


and ∴  ; a part


equal to the whole, which is abſurd:

∴ ——— and ———

do not biſect one another.

Q. E. D.



IF *two circles*  *interſect, they have not the ſame centre.*

Suppoſe it poſſible that two interſecting circles have a common centre; from ſuch ſuppoſed centre draw —— to the interſecting point, and ——····· meeting the circumferences of the circles.

Then —— = —— (B. 1. def. 15.)

and —— = ——····· (B. 1. def. 15.)


∴ —— = ——·····; a part equal to the whole, which is abſurd:

∴ circles ſuppoſed to interſect in any point cannot have the ſame centre.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *two circles*  *touch one another internally, they have not the ſame centre.*



For, if it be poſſible, let both circles have the ſame centre; from ſuch a ſuppoſed centre draw ----- cutting both circles, and ——— to the point of contact.

Then ——— = ----- (B. 1. def. 15.)

and ——— = ----- (B. 1. def. 15.)


$\therefore$ ----- = -----; a part equal to the whole, which is abſurd;

therefore the aſſumed point is not the centre of both circles; and in the ſame manner it can be demonſtrated that no other point is.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

FIGURE I.



IF *from any point within a circle* ○ *which is not the centre, lines* { *are drawn to the circumference; the greateſt of thoſe lines is that* (▬···) *which paſſes through the centre, and the leaſt is the remaining part* (▬) *of the diameter.*

Of the others, that (▬) *which is nearer to the line paſſing through the centre, is greater than that* (▬) *which is more remote.*

FIGURE II.




Fig. 2. *The two lines* (▬·· *and* ▬) *which make equal angles with that paſſing through the centre, on oppoſite ſides of it, are equal to each other; and there cannot be drawn a third line equal to them, from the ſame point to the circumference.*

FIGURE I.

To the centre of the circle draw ▬ and ▬; then ▬ = ▬ (B. 1. def. 15.)

▬ = ▬ + ▬ ⊏ ▬ (B. 1. pr. 20.)

in like manner ▬ may be ſhewn to be greater than ▬, or any other line drawn from the ſame point to the circumference. Again, by (B. 1. pr. 20.)

▬ + ▬ ⊏ ▬ = ▬ + ▬,

take ▬ from both; ∴ ▬ ⊏ ▬ (ax.),

and in like manner it may be ſhewn that ▬ is leſs

Harvard University - Houghton Library / pga_typ_805_47_3730

than any other line drawn from the ſame point to the cir-

cumference. Again, in  and  ,

 common, , and









∴  (B. 1. pr. 24.) and
may in like manner be proved greater than any other line drawn from the ſame point to the circumference more remote from  .

FIGURE II.



If  then  , if not
take  draw  , then
in  and  , common,

 and 

∴  (B. 1. pr. 4.)

∴ 


a part equal to the whole, which is abſurd:

∴  ; and no other line is equal to
 drawn from the ſame point to the circumference; for if it were nearer to the one paſſing through the centre it would be greater, and if it were more remote it would be leſs.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

The original text of this propoſition is here divided into three parts.



I.

IF *from a point without a circle, ſtraight lines* { —— ------ — —— — &c. } *are drawn to the circumference; of thoſe falling upon the concave circumference the greateſt is that* (——----) *which paſſes through the centre, and the line* (——) *which is nearer the greateſt is greater than that* (——) *which is more remote.*

Draw ..... and ..... to the centre.

Then, ——---- which paſſes through the centre, is greateſt; for ſince ------- = ------- , if —— be added to both, ——---- = —— + ------; but ⊏ —— (B. 1. pr. 20.) ∴ ——---- is greater than any other line drawn from the ſame point to the concave circumference.

Again in  and  , ... = ...,

Harvard University - Houghton Library / pga_typ_805_47_3730


and ——— common, but  ,

∴ ——— ⊏ ——— (B. 1. pr. 24.);

and in like manner ——— may be ſhewn ⊏ than any other line more remote from ———.

II.

Of thoſe lines falling on the convex circumference the leaſt is that (----------) *which being produced would paſs through the centre, and the line which is nearer to the leaſt is leſs than that which is more remote.*



For, ſince ——— + ····· ⊏ ———— (B. 1. pr. 20.)

and ——— = ———,

∴ ------- ⊏ ---------- (ax. 5.)

And again, ſince ——— + ····· ⊏


——— + ····· (B. 1. pr. 21.),

and ——— = ———,

∴ ····· ⊐ -----. And ſo of others.

III.

Alſo the lines making equal angles with that which paſſes through the centre are equal, whether falling on the concave or convex circumference; and no third line can be drawn equal to them from the ſame point to the circumference.



For if ····——— ⊏ ·····, but making ◢ = ◣;

make ------ = -----, and draw ····——.

Harvard University - Houghton Library / pga_typ_805_47_3730

Then in  and  we have ▬▬ = ▬▬,

and ▬▬ common, and alſo ◣ = ◢,

∴ ▬▬ = ▬▬ (B. 1. pr. 4.);

but ▬▬ = ▬▬;

∴ ▬▬ = ▬▬, which is abſurd.

∴ ▬▬ is not = ▬▬, nor to any part of ▬▬, ∴ ▬▬ is not ⊏ ▬▬.


Neither is ▬▬ ⊏ ▬▬, they are

∴ = to each other.

And any other line drawn from the ſame point to the circumference muſt lie at the ſame ſide with one of theſe lines, and be more or leſs remote than it from the line paſſing through the centre, and cannot therefore be equal to it.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *a point be taken within a circle* ○, *from which more than two equal ſtraight lines* (┈┈, ——, ———) *can be drawn to the circumference, that point muſt be the centre of the circle.*

For, if it be ſuppoſed that the point ⅄ in which more than two equal ſtraight lines meet is not the centre, ſome other point ——·· muſt be; join theſe two points by ———, and produce it both ways to the circumference.

Then ſince more than two equal ſtraight lines are drawn from a point which is not the centre, to the circumference, two of them at leaſt muſt lie at the ſame ſide of the diameter ——·····; and ſince from a point ⅄, which is not the centre, ſtraight lines are drawn to the circumference; the greateſt is ——┈┈, which paſſes through the centre: and ——— which is nearer to ——┈┈, ⊏ ——— which is more remote (B. 3. pr. 8.);

but ——— = ——— (hyp.) which is abſurd.

The ſame may be demonſtrated of any other point, different from ⅄, which muſt be the centre of the circle.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



ONE *circle* ○ *cannot interſect another* ○ *in more points than two.*

For, if it be poſſible, let it interſect in three points;

from the centre of ○ draw ——, ——

and —— to the points of interſection;

∴ —— = —— = ——

(B. 1. def. 15.),

but as the circles interſect, they have not the ſame centre (B. 3. pr. 5.):


∴ the aſſumed point is not the centre of ○, and

∴ as ——, —— and —— are drawn from a point not the centre, they are not equal (B. 3. prs. 7, 8); but it was ſhewn before that they were equal, which is abſurd; the circles therefore do not interſect in three points.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *two circles* ○ *and* ○ *touch one another internally, the right line joining their centres, being produced, ſhall paſs through a point of contact.*



For, if it be poſſible, let ——— join their centres, and produce it both ways; from a point of contact draw ——— to the centre of ○, and from the ſame point of contact draw ▪▪▪▪▪▪▪▪▪ to the centre of ○.



Becauſe in ; ——— + ——— ⊏ ▪▪▪▪▪▪▪▪▪

(B. 1. pr. 20.),

and ▪▪▪▪▪▪▪▪▪ = ——·——·—— as they are radii of ○,

but —— + —— ⊏ ——; take

away —— which is common,

and —— ⊏ ——;

but —— = ——,

becauſe they are radii of ◯,


and ∴ —— ⊏ —— a part greater than the whole, which is abſurd.

The centres are not therefore ſo placed, that a line joining them can paſs through any point but a point of contact.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *two circles* ○ *and* ○ *touch one another externally, the ſtraight line* —— *joining their centres, paſſes through the point of contact.*



If it be poſſible, let —— join the centres, and not paſs through a point of contact; then from a point of contact draw ------ and —— to the centres.

Becauſe ------ + —— ⊏ ——
(B. 1. pr. 20.),

and —— = ------ (B. 1. def. 15.),

and —— = —— (B. 1. def. 15.),

∴ — + — ⊏ ——, a part greater than the whole, which is abſurd.

The centres are not therefore ſo placed, that the line joining them can paſs through any point but the point of contact.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

FIGURE I.





FIGURE II.



ONE *circle cannot touch another, either externally or internally, in more points than one.*

FIGURE III.





Fig. 1. For, if it be poſſible, let  and  touch one another internally in two points; draw  joining their centres, and produce it until it paſs through one of the points of contact (B. 3. pr. 11.);


draw and  ,

But  (B. 1. def. 15.),

∴ if be added to both,

;

but (B. 1. def. 15.),

and ∴  ; but


 (B. 1. pr. 20.),

which is abſurd.


Harvard University - Houghton Library / pga_typ_805_47_3730

Fig. 2. But if the points of contact be the extremities of the right line joining the centres, this ſtraight line muſt be biſected in two different points for the two centres; becauſe it is the diameter of both circles, which is abſurd.

Fig. 3. Next, if it be poſſible, let  and  touch externally in two points; draw  joining the centres of the circles, and paſſing through one of the points of contact, and draw and .

 (B. 1. def. 15.);

and  (B. 1. def. 15.):

 but


 (B. 1. pr. 20.),

which is abſurd.

There is therefore no caſe in which two circles can touch one another in two points.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



EQUAL *ſtraight lines* ( ) *inſcribed in a circle are equally diſtant from the centre; and alſo, ſtraight lines equally diſtant from the centre are equal.*

From the centre of draw ⊥ to and ⊥ , join and .

Then = half (B. 3. pr. 3.)

and = ½ (B. 3. pr. 3.)

ſince = (hyp.)

∴ = ,

and = (B. 1. def. 15.)

∴ ² = ²;

but ſince is a right angle

² = ² + ² (B. 1. pr. 47.)

and ² = ² + ² for the ſame reaſon,

∴ ² + ² = ² + ²

Harvard University - Houghton Library / pga_typ_805_47_3730

$\therefore$ ----------^{2} = ----------^{2},

$\therefore$ ---------- = ----------.

Alſo, if the lines ——····· and ——···· be equally diſtant from the centre; that is to ſay, if the perpendiculars ---------- and ---------- be given equal, then

——····· = ——····.

For, as in the preceding caſe,

----------^{2} + ——^{2} = ——^{2} + ------^{2};

but ----------^{2} = ----------^{2}:


$\therefore$ ——^{2} = ——^{2}, and the doubles of theſe

——···· and ——····· are alſo equal.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

FIGURE I.



THE *diameter is the greateſt ſtraight line in a circle: and, of all others, that which is neareſt to the centre is greater than the more remote.*

FIGURE I.

The diameter —— is ⊏ any line ——.

For draw —— and ·····.

Then ····· = ——

and —— = ——,

∴ —— + ····· = ——

but —— + ····· ⊏ —— (B. 1. pr. 20.)

∴ —— ⊏ ——.

Again, the line which is nearer the centre is greater than the one more remote.

Firſt, let the given lines be —— and ·····,

which are at the ſame ſide of the centre and do not interſect;

draw { ····· , —— , ····· , ····· .

Harvard University - Houghton Library / pga_typ_805_47_3730

In  and  ,

——— and ------- = - - - - - - - and ····· ;

but  ⊏  ,

∴ ——— ⊏ - - - - - - (B. 1. pr. 24.)


FIGURE II.

Let the given lines be ——— and ——— which either are at different ſides of the centre, or interſect; from the centre draw ····· and - - - - - - ⊥ ——— and ———,

make ····· = ····, and

draw ——— ⊥ ····.

FIGURE II.




Since ——— and ——— are equally diſtant from the centre, ——— = ——— (B. 3. pr. 14.);

but ——— ⊏ ——— (Pt. 1. B. 3. pr. 15.),

∴ ——— ⊏ ———.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



THE *ſtraight line* ——— *drawn from the extremity of the diameter* ——— *of a circle perpendicular to it falls without the circle.*

And if any ſtraight line - - - - - *be drawn from a point within that perpendicular to the point of contact, it cuts the circle.*

PART I

If it be poſſible, let ———, which meets the circle again, be ⊥ ———, and draw ———.

Then, becauſe ——— = ———,

◢ = ◥ (B. 1. pr. 5.),


and ∴ each of these angles is acute. (B. 1. pr. 17.)

but ◢ = ◺ (hyp.), which is abſurd, therefore ——— drawn ⊥ ——— does not meet the circle again.

Harvard University - Houghton Library / pga_typ_805_47_3730

PART II.

Let ——— be ⊥ ——— and let - - - - - be drawn from a point ⋯ between ——— and the circle, which, if it be poſſible, does not cut the circle.



Becauſe = ,

∴ is an acute angle; ſuppoſe ⋯ ⊥ - - - - - , drawn from the centre of the circle, it muſt fall at the ſide of the acute angle.

∴ which is ſuppoſed to be a right angle, is ⊏ ,


∴ ——— ⊏ ⋯⋯;

but ⋯⋯ = ———,

and ∴ - - - - - ⊏ ⋯⋯⋯, a part greater than the whole, which is abſurd. Therefore the point does not fall outſide the circle, and therefore the ſtraight line ⋯⋯ cuts the circle.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730





TO *draw a tangent to a given circle* ◯ *from a given point, either in or outside of its circumference.*


If the given point be in the circumference, as at ....., it is plain that the straight line —— ⊥ ---------- the radius, will be the required tangent (B. 3. pr. 16.) But if the given point ⌝ be outside of the circumference, draw ·····—— from it to the centre, cutting ◯; and draw ---------- ⊥ ----------, describe ◯ concentric with ◯ radius = ·····—, then —— will be the tangent required.


Harvard University - Houghton Library / pga_typ_805_47_3730

For in  and 

 common,


and  ,

∴ (B. 1. pr. 4.)  a right angle,

∴  is a tangent to  .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *a right line* ..... *be a tangent to a circle, the ſtraight line* —— *drawn from the centre to the point of contact, is perpendicular to it.*

For, if it be poſſible,

let —··— be ⊥ ·····,

then becauſe ◺ = ◝,

◔ is acute (B. 1. pr. 17.)

∴ —— ⊏ —·—·—

(B. 1. pr. 19.);

but —— = ——,


and ∴ —·— ⊏ —·—·—, a part greater than the whole, which is abſurd.

∴ —·— is not ⊥ ·····; and in the ſame manner it can be demonſtrated, that no other line except —— is perpendicular to ·····.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *a ſtraight line* ——— *be a tangent to a circle, the ſtraight line* ———, *drawn perpendicular to it from point of the contact, paſſes through the centre of the circle.*



For, if it be poſſible, let the centre be without ———, and draw ..... from the ſuppoſed centre to the point of contact.

Becauſe ..... ⊥ ———
(B. 3. pr. 18.)

∴ ◜ = ◺, a right angle;

but ◜ = ◺ (hyp.), and ∴ ◜ = ◜,
a part equal to the whole, which is abſurd.

Therefore the aſſumed point is not the centre; and in the ſame manner it can be demonſtrated, that no other point without ——— is the centre.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

FIGURE I



THE *angle at the centre of a circle, is double the angle at the circumference, when they have the ſame part of the circumference for their baſe.*

FIGURE I.

Let the centre of the circle be on ——·····

a ſide of △.

Becauſe —— = ——,

△ = ◥ (B. 1. pr. 5.).

But ◓ = △ + ◥,

or ◓ = twice △ (B. 1. pr. 32).

FIGURE II.





FIGURE II.

Let the centre be within ◓, the angle at the circumference; draw —— from the angular point through the centre of the circle;


then ◥ = ◥, and △ = ◺,

becauſe of the equality of the ſides (B. 1. pr. 5).

Harvard University - Houghton Library / pga_typ_805_47_3730

Hence  twice .

But  , and

 ,

 twice .

FIGURE III.

Let the centre be without  and draw , the diameter.


Becauſe  twice  ; and

 twice  (caſe 1.);

 twice  .


Q. E. D.

FIGURE III.



Harvard University - Houghton Library / pga_typ_805_47_3730

FIGURE I.



THE *angles* (▲ , ▲) *in the ſame ſegment of a circle are equal.*

FIGURE I.

Let the ſegment be greater than a ſemicircle, and draw ——— and ——— to the centre.

◗ = twice ▲ or twice = ▲

(B. 3. pr. 20.);

∴ ▲ = ▲.

FIGURE II.




FIGURE II.

Let the ſegment be a ſemicircle, or leſs than a ſemicircle, draw ——— the diameter, alſo draw ———.


◗ = ▲ and ◥ = ◣ (caſe 1.)


∴ ◗ = ◣.


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

THE *oppoſite angles*  *and*  *and*  *of any quadrilateral figure inſcribed in a circle, are together equal to two right angles.*



Draw —— and —— the diagonals; and becauſe angles in the ſame ſegment are equal  ,

and  ;

add  to both.

∴ 

two right angles (B. 1. pr. 32.). In like manner it may be ſhown that,


 .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



UPON *the ſame ſtraight line, and upon the ſame ſide of it, two ſimilar ſegments of circles cannot be conſtructed which do not coincide.*


For if it be poſſible, let two ſimilar ſegments  and  be conſtructed;

draw any right line ——— cutting both the ſegments,

draw ——— and ——— .

Becauſe the ſegments are ſimilar,

 (B. 3. def. 10.),

but  (B. 1. pr. 16.)

which is abſurd: therefore no point in either of the ſegments falls without the other, and therefore the ſegments coincide.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

SIMILAR *ſegments*  *and*  , *of circles upon equal ſtraight lines* (—— *and* ——) *are each equal to the other.*






For, if  be ſo applied to  ,
that —— may fall on ——, the extremities of
—— may be on the extremities —— and
 at the ſame ſide as  ;
becauſe —— = ——,
—— muſt wholly coincide with ——;
and the ſimilar ſegments being then upon the ſame
ſtraight line and at the ſame ſide of it, muſt
alſo coincide (B. 3. pr. 23.), and
are therefore equal.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



A SEGMENT *of a circle being given, to deſcribe the circle of which it is the ſegment.*

From any point in the ſegment draw ——— and ——— biſect them, and from the points of biſection draw ——— ⊥ ———

and ——— ⊥ ———


where they meet is the centre of the circle.

Becauſe ——— terminated in the circle is biſected perpendicularly by ———, it paſſes through the centre (B. 3. pr. 1.), likewiſe ——— paſſes through the centre, therefore the centre is in the interſection of theſe perpendiculars.

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730

IN *equal circles* ○ *and* ○, *the arcs* ◡, ◡ *on which ſtand equal angles, whether at the centre or circumference, are equal.*



Firſt, let ◔ = ◔ at the centre,

draw ——— and - - - - - - - - .



Then ſince ○ = ○,

△ and △ have

——— = ——— = ------- = - - - - - - - ,

and ◔ = ◔,

∴ ——————— = - - - - - - - - (B. 1. pr. 4.).

But ▲ = ▲ (B. 3. pr. 20.);

∴ ⌒ and ⌒ are ſimilar (B. 3. def. 10.);

they are alſo equal (B. 3. pr. 24.)

Harvard University - Houghton Library / pga_typ_805_47_3730

If therefore the equal ſegments be taken from the equal circles, the remaining ſegments will be equal;

hence ⌓ = ⌓ (ax. 3.);

and ∴ ⌒ = ⌒.

But if the given equal angles be at the circumference, it is evident that the angles at the centre, being double of thoſe at the circumference, are alſo equal, and therefore the arcs on which they ſtand are equal.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IN *equal circles,* ○ ○ *an*



the angles ▲ *and* ▲ *which ſtand upon equal arches are equal, whether they be at the centres or at the circumferences.*

For if it be poſſible, let one of them ▲ be greater than the other ▲



and make

▲ = ▲

∴ ⌒ = ⌒ (B. 3. pr. 26.)

but ⌒ = ⌒ (hyp.)

∴ ⌒ = ⌒ a part equal


to the whole, which is abſurd; ∴ neither angle is greater than the other, and ∴ they are equal.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IN *equal circles* ○ *and* ○, *equal chords* ——, - - - - *cut off equal arches.*



From the centres of the equal circles,

draw ——, —— and - - - -, - - - -;

and becauſe ○ = ○

——, —— = - - - -, - - - -

alſo —— = - - - - (hyp.)

∴ ◔ = ◔

∴ ⌒ = ⌒ (B. 3. pr. 26.)

and ∴ ◠ = ◠ (ax. 3.)

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730

IN *equal circles*  *and*  

the chords ——— *and* - - - - - - - *which ſub-*

tend equal arcs are equal.

If the equal arcs be ſemicircles the propoſition is evident. But if not,



let ———, ———, and - - - - - - -, ... be drawn to the centres;

becauſe ⌒ = ⌒ (hyp.)

and ◔ = ◔ (B. 3. pr. 27.);


but ——— and ——— = - - - - - - - and .....

∴ ——— = ... (B. 1. pr. 4.);

but theſe are the chords ſubtending the equal arcs.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *bisect a given arc* .

Draw ——·····;

make —— = ·····,

draw —— ⊥ ——·····, and it bisects the arc.

Draw ····· and ——.

—— = ------ (const.),

—— is common,

and ◗ = ◔ (const.)

∴ ····· = —— (B. 1. pr. 4.)

⌒ = ⌒ (B. 3. pr. 28.),

and therefore the given arc is bisected.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IN *a circle the angle in a ſemicircle is a right angle, the angle in a ſegment greater than a ſemicircle is acute, and the angle in a ſegment leſs than a ſemicircle is obtuſe.*

FIGURE I.



FIGURE I.

The angle  in a ſemicircle is a right angle.

Draw ——— and ———

 (B. 1. pr. 5.)

 the half of two

right angles = a right angle. (B. 1. pr. 32.)

FIGURE II.



The angle in a ſegment greater than a ſemicircle is acute.

Draw ——— the diameter, and ———



∴ = a right angle



∴ is acute.

FIGURE II.





FIGURE III.




FIGURE III.

The angle  in a ſegment leſs than ſemicircle is obtuſe.

Take in the oppoſite circumference any point, to which draw and .

Becauſe 

(B. 3. pr. 22.)


but  (part 2.),

∴  is obtuſe.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *a right line* —— *be a tangent to a circle, and from the point of contact a right line* —— *be drawn cutting the circle, the angle* ◣ *made by this line with the tangent is equal to the angle* ◤ *in the alternate ſegment of the circle.*



If the chord ſhould paſs through the centre, it is evident the angles are equal, for each of them is a right angle. (B. 3. prs. 16, 31.)

But if not, draw —— ⊥ —— from the point of contact, it muſt paſs through the centre of the circle, (B. 3. pr. 19.)

∴ ◥ = ◮ (B. 3. pr. 31.)

◤ + ◥ = ◮ = ◥ (B. 1. pr. 32.)

∴ ◤ = ◣ (ax.).


Again ◮ = ◠◠ = ◤ + ◆

(B. 3. pr. 22.)

∴ ◮ = ◆, (ax.), which is the angle in the alternate ſegment.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



ON *a given ſtraight line* —— *to deſcribe a ſegment of a circle that ſhall contain an angle equal to a given angle* ◺, ⬠, ▲.

If the given angle be a right angle, biſect the given line, and deſcribe a ſemicircle on it, this will evidently contain a right angle. (B. 3. pr. 31.)

If the given angle be acute or obtuſe, make with the given line, at its extremity, ◢ = ◢, draw —— ⊥ —— and make ◥ = ◥, deſcribe ◯ with —— or —— as radius, for they are equal.


—— is a tangent to ◯ (B. 3. pr. 16.)

∴ —— divides the circle into two ſegments capable of containing angles equal to ◔ and ◢ which were made reſpectively equal to ⬠ and ▲ (B. 3. pr. 32.)


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *cut off from a given circle*  *a ſegment which ſhall contain an angle equal to a given angle*  .



Draw ——— (B. 3. pr. 17.), a tangent to the circle at any point; at the point of contact make

 = the given angle;

and  contains an angle = the given angle.

Becauſe ——— is a tangent,

and ——— cuts it, the


angle in  =  (B. 3. pr. 32.),

but  =  (conſt.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

FIGURE I.



IF *two chords* { — ····· / ---- — } *in a circle interſect each other, the rectangle contained by the ſegments of the one is equal to the rectangle contained by the ſegments of the other.*

FIGURE I.

If the given right lines paſs through the centre, they are biſected in the point of interſection, hence the rectangles under their ſegments are the ſquares of their halves, and are therefore equal.

FIGURE II.




FIGURE II.

Let ---·---- paſs through the centre, and ——····· not; draw ——— and ———.

Then ——— × ····· = ———^{2} — ·····^{2} (B. 2. pr. 6.),

or ——— × ····· = ·····^{2} — ·····^{2},

∴ ——— × ---—— = ·····—— × ---—— (B. 2. pr. 5.).

FIGURE III.




FIGURE III.

Let neither of the given lines paſs through the centre, draw through their interſection a diameter ·····——,

and ····· × —— = ——— × ····· (Part. 2.),

alſo ——···· × —— = ——— × ---—— (Part. 2.);

∴ ——— × ····· = ——— × ---——.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *from a point without a circle two ſtraight lines be drawn to it, one of which* ——— *is a tangent to the circle, and the other* — ····· — *cuts it; the rectangle under the whole cutting line* — ····· — *and the external ſegment* ——— *is equal to the ſquare of the tangent* ——— .

FIGURE I.




FIGURE I.

Let — ····· — paſs through the centre;

draw ——— from the centre to the point of contact;

———^{2} = ·····—^{2} minus ———^{2} (B. 1. pr. 47),


or ———^{2} = ·····—^{2} minus ·····^{2},

$\therefore$ ———^{2} = — ····· — × ——— (B. 2. pr. 6).

FIGURE II.

If ····— do not paſs through the centre, draw ····— and ····— .

FIGURE II.



Then ····— × —

= ———^{2} minus ····^{2}


(B. 2. pr. 6), that is,

····— × — = ———^{2} minus ———^{2},

$\therefore$ ····— × — = ———^{2} (B. 3. pr. 18).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *from a point outside of a circle two straight lines be drawn, the one* ▬·▬·▬ *cutting the circle, the other* ▬▬▬ *meeting it, and if the rectangle contained by the whole cutting line* ▬·····▬ *and its external segment* ▬▬▬▬ *be equal to the square of the line meeting the circle, the latter* ▬▬▬ *is a tangent to the circle.*

Draw from the given point ▬▬▬, a tangent to the circle, and draw from the centre ▬·▬, ▬▬▬, and ▬▬▬,

▬▬▬^{2} = ▬▬▬▬▬ × ▬▬▬▬ (B. 3. pr. 36.)

but ▬▬▬^{2} = ▬▬▬▬▬ × ▬▬▬▬ (hyp.),


$\therefore$ ▬▬▬^{2} = ▬▬▬^{2},


and $\therefore$ ▬▬▬ = ▬▬▬;


Then in  and 

▬▬▬ and ▬▬▬ = ▬▬▬ and ▬▬▬,

and ▬▬▬ is common,

$\therefore$  (B. 1. pr. 8.);

but  a right angle (B. 3. pr. 18.),

$\therefore$  a right angle,

and $\therefore$ ▬▬▬ is a tangent to the circle (B. 3. pr. 16.).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

BOOK IV.

DEFINITIONS.

I.

A RECTILINEAR figure is ſaid to be *inſcribed in* another, when all the angular points of the inſcribed figure are on the ſides of the figure in which it is ſaid to be inſcribed.



II.

A FIGURE is ſaid to be *deſcribed about* another figure, when all the ſides of the circumſcribed figure paſs through the angular points of the other figure.

III.

A RECTILINEAR figure is ſaid to be *inſcribed in* a circle, when the vertex of each angle of the figure is in the circumference of the circle.



IV.

A RECTILINEAR figure is ſaid to be *circumſcribed about* a circle, when each of its ſides is a tangent to the circle.




Harvard University - Houghton Library / pga_typ_805_47_3730



V.

A CIRCLE is ſaid to be *inſcribed in* a rectilinear figure, when each ſide of the figure is a tangent to the circle.

VI.



A CIRCLE is ſaid to be *circumſcribed about* a rectilinear figure, when the circumference paſſes through the vertex of each angle of the figure.



is circumſcribed.




VII.

A STRAIGHT line is ſaid to be *inſcribed in* a circle, when its extremities are in the circumference.

The Fourth Book of the Elements is devoted to the ſolution of problems, chiefly relating to the inſcription and circumſcription of regular polygons and circles.

A regular polygon is one whoſe angles and ſides are equal.

Harvard University - Houghton Library / pga_typ_805_47_3730



IN *a given circle* ○ *to place a ſtraight line, equal to a given ſtraight line* (——), *not greater than the diameter of the circle.*

Draw ·····—, the diameter of ○;

and if ·····— = ——, then

the problem is ſolved.

But if ·····— be not equal to ——,

·····— ⊏ —— (hyp.);

make ····· = —— (B. 1. pr. 3.) with

------ as radius,

deſcribe ○, cutting ○, and


draw ——, which is the line required.

For —— = ····· = ——

(B. 1. def. 15. conſt.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IN *a given circle* ○ *to inſcribe a triangle equiangular to a given triangle.*

To any point of the given circle draw ——, a tangent (B. 3. pr. 17.); and at the point of contact make ▲ = ▼ (B. 1. pr. 23.)

and in like manner ▲ = ▼, and draw ——.

Becauſe ▲ = ▼ (conſt.)

and ▲ = ◥ (B. 3. pr. 32.)

∴ ◥ = ▼; alſo

▽ = ▼ for the ſame reaſon.


∴ ▼ = ▼ (B. 1. pr. 32.),

and therefore the triangle inſcribed in the circle is equiangular to the given one.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730






ABOUT *a given circle* ○ *to circumſcribe a triangle equiangular to a given triangle.*

Produce any ſide ——, of the given triangle both ways; from the centre of the given circle draw ——, any radius.

Make ◔ = ◔ (B. 1. pr. 23.)


and ◕ = ◔.


At the extremities of the three radii, draw ——, —— and ----, tangents to the given circle. (B. 3. pr. 17.)

The four angles of  , taken together, are equal to four right angles. (B. 1. pr. 32.)


Harvard University - Houghton Library / pga_typ_805_47_3730

but  and  are right angles (conſt.)


$\therefore$  two right angles


but  (B. 1. pr. 13.)

and  (conſt.)

and $\therefore$  .

In the ſame manner it can be demonſtrated that

 ;



$\therefore$  (B. 1. pr. 32.)



and therefore the triangle circumſcribed about the given circle is equiangular to the given triangle.

Q. E. D.

IN *a given triangle*  *to in-ſcribe a circle.*



Biſect  and  (B. 1. pr. 9.) by ····· and ——; from the point where theſe lines meet draw - - - - - -, — — and - - - - reſpectively perpendicular to ———, —— and ——.

In  and 

 and ····· common, ∴ ····· = ····· (B. 1. pr. 4 and 26.)


In like manner, it may be ſhown alſo that ····· = ·····,

∴ - - - - - - - = ····· = ·····;

hence with any one of theſe lines as radius, deſcribe

 and it will paſs through the extremities of the other two; and the ſides of the given triangle, being perpendicular to the three radii at their extremities, touch the circle (B. 3. pr. 16.), which is therefore inſcribed in the given circle.


s Q. E. D.




TO *describe a circle about a given triangle.*

Make ——— = ····· and ——— = ····· (B. 1. pr. 10.)

From the points of bisection draw ——— and ····· ⊥ ——— and ——— respectively (B. 1. pr. 11.), and from their point of concourse draw ———, ····· and ——— and describe a circle with any one of them, and it will be the circle required.






In  and 

····· = ——— (const.),

——— common,

 (const.),

∴ ——— = ····· (B. 1. pr. 4.).

In like manner it may be shown that


——— = ·····.

∴ ····· = ——— = ———; and therefore a circle described from the concourse of these three lines with any one of them as a radius will circumscribe the given triangle.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IN *a given circle* ◯ *to inſcribe a ſquare.*



Draw the two diameters of the circle ⊥ to each other, and draw —, —, — and —

◇ is a ſquare.

For, ſince ◔ and ◕ are, each of them, in a ſemicircle, they are right angles (B. 3. pr. 31),

∴ — ∥ — (B. 1. pr. 28):

and in like manner — ∥ —.

And becauſe ◔ = ◔ (conſt.), and

┈┈ = ╍╍ = ┈┈ (B. 1. def. 15).

∴ — = — (B. 1. pr. 4);

and ſince the adjacent ſides and angles of the parallelogram ◇ are equal, they are all equal (B. 1. pr. 34);

and ∴ ◇, inſcribed in the given circle, is a ſquare.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



ABOUT *a given circle* ○ *to circumſcribe a ſquare.*

Draw two diameters of the given circle perpendicular to each other, and through their extremities draw ——, ——, ——, and —— tangents to the circle;

and □ is a ſquare.

◔ = ◺ a right angle, (B. 3. pr. 18.)

alſo ◕ = ◺ (conſt.),

∴ —— ‖ ····· ; in the ſame manner it can be demonſtrated that —— ‖ ····· , and alſo that —— and —— ‖ ····· ;

∴ □ is a parallelogram, and

becauſe ◔ = ◕ = ◕ = ◕ = ◕

they are all right angles (B. 1. pr. 34):

it is alſo evident that —— , —— , ——


and —— are equal.

∴ □ is a ſquare.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *inſcribe a circle in a given ſquare.*



Make ——— = ·····,

and ——— = ·····,

draw ·····— ‖ —·····,

and ▬▬▬ ‖ ·····

(B. 1. pr. 31.)

∴ ■ is a parallelogram;

and ſince —·····— = ·····— (hyp.)

——— = ·····

∴ ■ is equilateral (B. 1. pr. 34.)

In like manner, it can be ſhown that


■ = ■ are equilateral parallelograms;

∴ ▬▬▬ = ····· = ——— = ———,

and therefore if a circle be deſcribed from the concourſe of theſe lines with any one of them as radius, it will be inſcribed in the given ſquare. (B. 3. pr. 16.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *deſcribe a circle about a given ſquare*  .

Draw the diagonals ▬▪▬▪ and ▬▪▪▪▪ interſecting each other; then,


becauſe  and  have their ſides equal, and the baſe ▬▪▪▪▪ common to both,

 (B. 1. pr. 8),

or  is biſected: in like manner it can be ſhown

that  is biſected;

but  ,

hence  their halves,

∴ ▬▬ = ▬▬; (B. 1. pr. 6.)

and in like manner it can be proved that


▬▬ = ▬▬ = ▪▪▪▪▪▪▪▪▪▪ = ▪▪▪▪▪▪▪▪▪▪.

If from the confluence of theſe lines with any one of them as radius, a circle be deſcribed, it will circumſcribe the given ſquare.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *conſtruct an iſoſceles triangle, in which each of the angles at the baſe ſhall be double of the vertical angle.*



Take any ſtraight line ——····· and divide it ſo that

—— · ····· = ——^{2}

(B. 2. pr. 11.)

With ——····· as radius, deſcribe ◯ and place in it from the extremity of the radius, —— = ——, (B. 4. pr. 1); draw ——.

Then ◺ is the required triangle.

For, draw —— and deſcribe

◯ about ◿ (B. 4. pr. 5.)


Since —— · ····· = ——^{2} = ——^{2},


∴ —— is a tangent to ◯ (B. 3. pr. 37.)

∴ ▲ = ◿ (B. 3. pr. 32),


Harvard University - Houghton Library / pga_typ_805_47_3730

add  to each,



but  (B. 1. pr. 5):


since  (B. 1. pr. 5.)


consequently 


(B. 1. pr. 32.)










 twice ; and consequently each angle at the base is double of the vertical angle.

Q. E. D.



IN a given circle  to inſcribe an equilateral and equiangular pentagon.

Conſtruct an iſoſceles triangle, in which each of the angles at the baſe ſhall be double of the angle at the vertex, and inſcribe in the given circle a triangle  equiangular to it; (B. 4. pr. 2.)

Biſect  and  (B. 1. pr. 9.)

draw ——, ——, ——— and - - - - - - -.

Becauſe each of the angles



are equal,

the arcs upon which they ſtand are equal, (B. 3. pr. 26.)


and ∴ ——, ——, ——, ——— and

- - - - - - - - - which ſubtend theſe arcs are equal (B. 3. pr. 29.)

and ∴ the pentagon is equilateral, it is alſo equiangular, as each of its angles ſtand upon equal arcs. (B. 3. pr. 27).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *deſcribe an equilateral and equiangular pentagon about a given circle* ○.

Draw five tangents through the vertices of the angles of any regular pentagon inſcribed in the given circle ○ (B. 3. pr. 17).

Theſe five tangents will form the required pentagon.

Draw { }. In ▽ and ◺

——— = ——— (B. 1. pr. 47),

- - - - - - - - - = · · · · · · · · ·, and ——— common;

∴ ▽ = ◡ and ◥ = ◢ (B. 1. pr. 8.)

∴ ◔ = twice ◡, and ◔ = twice ◢;


In the ſame manner it can be demonſtrated that


◔ = twice ◡, and ◔ = twice ◢;

but ◔ = ◔ (B. 3. pr. 27),


Harvard University - Houghton Library / pga_typ_805_47_3730


∴ their halves  common;






In the ſame manner it can be demonſtrated that the other ſides are equal, and therefore the pentagon is equilateral, it is alſo equiangular, for





demonſtrated that the other angles of the deſcribed pentagon are equal.

Q. E. D




TO *inſcribe a circle in a given equiangular and equilateral pentagon.*

Let  be a given equiangular and equilateral pentagon; it is required to inſcribe a circle in it.

Make  =  , and  = 
(B. 1. pr. 9.)

Draw  , , , , &c.

Becauſe = ,  =  ,
and common to the two triangles

 and  ;

∴ = and  =  (B. 1. pr. 4.)



And becauſe  =  = twice 



∴ = twice  , hence  is biſected by .



In like manner it may be demonſtrated that  is biſected by  , and that the remaining angle of the polygon is biſected in a ſimilar manner.

Harvard University - Houghton Library / pga_typ_805_47_3730

Draw ——— , - - - - - - - - , &c. perpendicular to the ſides of the pentagon.

Then in the two triangles  and 

we have  =  , (conſt.), ——— common,

and  =  = a right angle;


∴ ——— = - - - - - - - - . (B. 1. pr. 26.)

In the ſame way it may be ſhown that the five perpendiculars on the ſides of the pentagon are equal to one another.


Deſcribe with any one of the perpendiculars as radius, and it will be the inſcribed circle required. For if it does not touch the ſides of the pentagon, but cut them, then a line drawn from the extremity at right angles to the diameter of a circle will fall within the circle, which has been ſhown to be abſurd. (B. 3. pr. 16.)


Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730




TO *deſcribe a circle about a given equilateral and equiangular pentagon.*

Biſect  and by ····· and ·····, and from the point of ſection, draw ——, ·····, and ——.

 ,

 , ∴ ····· = ····· (B. 1. pr. 6);

and ſince in  and  ,

—— = ——, and ····· common,

alſo  = ;

∴ —— = ····· (B. 1. pr. 4).

In like manner it may be proved that

····· = —— = ——, and


therefore ····· = —— = ····· = ····· = ——:

Therefore if a circle be deſcribed from the point where theſe five lines meet, with any one of them as a radius, it will circumſcribe the given pentagon.



Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730


TO *inſcribe an equilateral and equiangular hexagon in a given circle* ○.



From any point in the circumference of the given circle deſcribe ○ paſſing through its centre, and draw the diameters ——, —— and ——; draw - - - - - -, · · · · · ·, · · · · · ·, &c. and the required hexagon is inſcribed in the given circle.

Since —— paſſes through the centres of the circles,  and  are equilateral triangles, hence  = one-third of two right angles; (B. 1. pr. 32) but  = 


(B. 1. pr. 13);

∴  = one-third of 

(B. 1. pr. 32), and the angles vertically oppoſite to theſe are all equal to one another (B. 1. pr. 15), and ſtand on equal arches (B. 3. pr. 26), which are ſubtended by equal chords (B. 3. pr. 29); and ſince each of the angles of the hexagon is double of the angle of an equilateral triangle, it is alſo equiangular.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *inſcribe an equilateral and equiangular quindecagon in a given circle.*

Let ——— and ——— be the ſides of an equilateral pentagon inſcribed in the given circle, and ——— the ſide of an inscribed equilateral triangle.

The arc ſubtended by ——— and ——— } = $\frac{2}{5}$ = $\frac{6}{15}$ { of the whole circumference.

The arc ſubtended by ——— } = $\frac{1}{3}$ = $\frac{5}{15}$ { of the whole circumference.

Their difference = $\frac{1}{15}$

∴ the arc ſubtended by ▪▪▪▪▪▪▪▪▪▪ = $\frac{1}{15}$ difference of the whole circumference.

Hence if ſtraight lines equal to ▪▪▪▪▪▪▪▪▪▪ be placed in the circle (B. 4. pr. 1), an equilateral and equiangular quindecagon will be thus inſcribed in the circle.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

BOOK V.

DEFINITIONS.

I.

LESS magnitude is ſaid to be an aliquot part or ſubmultiple of a greater magnitude, when the leſs meaſures the greater; that is, when the leſs is contained a certain number of times exactly in the greater.

II.

A GREATER magnitude is ſaid to be a multiple of a leſs, when the greater is meaſured by the leſs; that is, when the greater contains the leſs a certain number of times exactly.

III.

RATIO is the relation which one quantity bears to another of the ſame kind, with reſpect to magnitude.

IV.

MAGNITUDES are ſaid to have a ratio to one another, when they are of the ſame kind; and the one which is not the greater can be multiplied ſo as to exceed the other.

The other definitions will be given throughout the book where their aid is firſt required.

Harvard University - Houghton Library / pga_typ_805_47_3730

AXIOMS.

I.

EQUIMULTIPLES or equiſubmultiples of the ſame, or of equal magnitudes, are equal.

If A = B, then
twice A = twice B, that is,
2 A = 2 B;
3 A = 3 B;
4 A = 4 B;
&c. &c.
and $\frac{1}{2}$ of A = $\frac{1}{2}$ of B;
$\frac{1}{3}$ of A = $\frac{1}{3}$ of B;
&c. &c.

II.

A MULTIPLE of a greater magnitude is greater than the ſame multiple of a leſs.

Let A ⊏ B, then
2 A ⊏ 2 B;
3 A ⊏ 3 B;
4 A ⊏ 4 B;
&c. &c.

III.

THAT magnitude, of which a multiple is greater than the ſame multiple of another, is greater than the other.

Let 2 A ⊏ 2 B, then
A ⊏ B;
or, let 3 A ⊏ 3 B, then
A ⊏ B;
or, let *m* A ⊏ *m* B, then
A ⊏ B.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *any number of magnitudes be equimultiples of as many others, each of each: what multiple soever any one of the firſt is of its part, the ſame multiple ſhall of the firſt magnitudes taken together be of all the others taken together.*

Let  be the ſame multiple of  ,

that  is of .

that  is of .

Then is evident that

 is the ſame multiple of 

which that  is of  ;

becauſe there are as many magnitudes

in  = 

as there are in  =  .

The ſame demonſtration holds in any number of magnitudes, which has here been applied to three.

∴ If any number of magnitudes, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730


IF *the firſt magnitude be the ſame multiple of the ſecond that the third is of the fourth, and the fifth the ſame multiple of the ſecond that the ſixth is of the fourth, then ſhall the firſt, together with the fifth, be the ſame multiple of the ſecond that the third, together with the ſixth, is of the fourth.*

Let ●●●, the firſt, be the ſame multiple of ●, the ſecond, that ◇◇◇, the third, is of ◇, the fourth; and let ●●●●, the fifth, be the ſame multiple of ●, the ſecond, that ◇◇◇◇, the ſixth, is of ◇, the fourth.


Then it is evident, that $\left\{\begin{matrix}●●● \\ ●●●●\end{matrix}\right\}$, the firſt and fifth together, is the ſame multiple of ●, the ſecond, that $\left\{\begin{matrix}◇◇◇ \\ ◇◇◇◇\end{matrix}\right\}$, the third and ſixth together, is of the ſame multiple of ◇, the fourth; becauſe there are as many magnitudes in $\left\{\begin{matrix}●●● \\ ●●●●\end{matrix}\right\}$ = ● as there are in $\left\{\begin{matrix}◇◇◇ \\ ◇◇◇◇\end{matrix}\right\}$ = ◇.

∴ If the firſt magnitude, &c.

IF *the firſt of four magnitudes be the ſame multiple of the ſecond that the third is of the fourth, and if any equimultiples whatever of the firſt and third be taken, thoſe ſhall be equimultiples; one of the ſecond, and the other of the fourth.*





Then it is evident,



which  is of  ;

The Fourth.

becauſe  contains  contains 

as many times as

 contains  contains  .

The ſame reaſoning is applicable in all caſes.


∴ If the firſt four, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

DEFINITION V.

Four magnitudes, ●, ■, ◆, ▼, are ſaid to be proportionals when every equimultiple of the firſt and third be taken, and every equimultiple of the ſecond and fourth, as,

of the firſt




&c.

of the third




&c.

of the ſecond



&c.


of the fourth




&c.

Then taking every pair of equimultiples of the firſt and third, and every pair of equimultiples of the ſecond and fourth,

If





then will



Harvard University - Houghton Library / pga_typ_805_47_3730


That is, if twice the firſt be greater, equal, or leſs than twice the ſecond, twice the third will be greater, equal, or leſs than twice the fourth; or, if twice the firſt be greater, equal, or leſs than three times the ſecond, twice the third will be greater, equal, or leſs than three times the fourth, and ſo on, as above expreſſed.






In other terms, if three times the firſt be greater, equal, or leſs than twice the ſecond, three times the third will be greater, equal, or leſs than twice the fourth; or, if three times the firſt be greater, equal, or leſs than three times the ſecond, then will three times the third be greater, equal, or leſs than three times the fourth; or if three times the firſt be greater, equal, or leſs than four times the ſecond, then will three times the third be greater, equal, or leſs than four times the fourth, and ſo on. Again,

Harvard University - Houghton Library / pga_typ_805_47_3730






And so on, with any other equimultiples of the four magnitudes, taken in the ſame manner.

Euclid expreſſes this definition as follows:—

The firſt of four magnitudes is ſaid to have the ſame ratio to the ſecond, which the third has to the fourth, when any equimultiples whatſoever of the firſt and third being taken, and any equimultiples whatſoever of the ſecond and fourth; if the multiple of the firſt be leſs than that of the ſecond, the multiple of the third is alſo leſs than that of the fourth; or, if the multiple of the firſt be equal to that of the ſecond, the multiple of the third is alſo equal to that of the fourth; or, ir the multiple of the firſt be greater than that of the ſecond, the multiple of the third is alſo greater than that of the fourth.

In future we ſhall expreſs this definition generally, thus:



x

Then we infer that ●, the firſt, has the ſame ratio to ■, the ſecond, which ◆, the third, has to ▼ the fourth: expreſſed in the ſucceeding demonſtrations thus:

● : ■ :: ◆ : ▼;

or thus, ● : ■ = ◆ : ▼;

or thus, $\frac{●}{■} = \frac{◆}{▼}$: and is read,

"as ● is to ■, so is ◆ to ▼."

And if ● : ■ :: ◆ : ▼ we ſhall infer if

M ● ⊏, = or ⊐ *m* ■, then will

M ◆ ⊏, = or ⊐ *m* ▼.

That is, if the firſt be to the ſecond, as the third is to the fourth; then if M times the firſt be greater than, equal to, or leſs than *m* times the ſecond, then ſhall M times the third be greater than, equal to, or leſs than *m* times the fourth, in which M and *m* are not to be conſidered particular multiples, but every pair of multiples whatever; nor are ſuch marks as ●, ▼, ■, &c. to be conſidered any more than repreſentatives of geometrical magnitudes.

The ſtudent ſhould thoroughly underſtand this definition before proceeding further.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *the firſt of four magnitudes have the ſame ratio to the ſecond, which the third has to the fourth, then any equimultiples whatever of the firſt and third ſhall have the ſame ratio to any equimultiples of the ſecond and fourth; viz., the equimultiple of the firſt ſhall have the ſame ratio to that of the ſecond, which the equimultiple of the third has to that of the fourth.*

Let ● : ■ :: ◆ : ⛊, then 3 ● : 2 ■ :: 3 ◆ : 2 ⛊, every equimultiple of 3 ● and 3 ◆ are equimultiples of ● and ◆, and every equimultiple of 2 ■ and 2 ⛊, are equimultiples of ■ and ⛊ (B. 5, pr. 3.)

That is, M times 3 ● and M times 3 ◆ are equimultiples of ● and ◆, and *m* times 2 ■ and *m* 2 ⛊ are equimultiples of 2 ■ and 2 ⛊; but ● : ■ :: ◆ : ⛊ (hyp); ∴ if M 3 ● ⊏, =, or ⊐ *m* 2 ■, then

M 3 ◆ ⊏, =, or ⊐ *m* 2 ⛊ (def. 5.)


and therefore 3 ● : 2 ■ :: 3 ◆ : 2 ⛊ (def. 5.)

The ſame reaſoning holds good if any other equimultiple of the firſt and third be taken, any other equimultiple of the ſecond and fourth.

∴ If the firſt four magnitudes, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *one magnitude be the ſame multiple of another, which a magnitude taken from the firſt is of a magnitude taken from the other, the remainder ſhall be the ſame multiple of the remainder, that the whole is of the whole.*



∴ If one magnitude, &c.

IF *two magnitudes be equimultiples of two others, and if equimultiples of these be taken from the first two, the remainders are either equal to these others, or equimultiples of them.*

Let ◇◇ = M′ ■ ; and ⊔⊔ = M′ ▲ ;

then ◇◇ minus m' ■ =

M′ ■ minus m' ■ = (M′ minus m') ■,

and ⊔⊔ minus m' ▲ = M′ ▲ minus m' ▲ =

(M′ minus m') ▲ .

Hence, (M′ minus m') ■ and (M′ minus m') ▲ are equimultiples of ■ and ▲ , and equal to ■ and ▲ , when M′ minus m' = 1.

∴ If two magnitudes be equimultiples, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *the first of the four magnitudes has the same ratio to the second which the third has to the fourth, then if the first be greater than the second, the third is also greater than the fourth; and if equal, equal; if less, less.*

Let ● : ■ :: ⛊ : ◆ ; therefore, by the fifth definition, if ● ● ⊏ ■ ■, then will ⛊ ⛊ ⊏ ◆ ◆ ;

but if ● ⊏ ■, then ● ● ⊏ ■ ■

and ⛊ ⛊ ⊏ ◆ ◆ ,

and ∴ ⛊ ⊏ ◆ .

Similarly, if ● ═, or ⊐ ■, then will ⛊ ═,

or ⊐ ◆ .

∴ If the first of four, &c.

DEFINITION XIV.

GEOMETRICIANS make use of the technical term "Invertendo," by inversion, when there are four proportionals, and it is inferred, that the second is to the first as the fourth to the third.

Let A : B :: C : D, then, by "invertendo" it is inferred B : A :: D : C.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *four magnitudes are proportionals, they are proportionals alſo when taken inverſely.*

Let ▼ : ∪ :: ■ : ◆,

then, inverſely, ∪ : ▼ :: ◆ : ■.

If M ▼ ⊐ *m* ∪, then M ■ ⊐ *m* ◆
by the fifth definition.

Let M ▼ ⊐ *m* ∪, that is, *m* ∪ ⊏ M ▼,

∴ M ■ ⊐ *m* ◆, or, *m* ◆ ⊏ M ■;

∴ if *m* ∪ ⊏ M ▼, then will *m* ◆ ⊏ M ■.

In the ſame manner it may be ſhown,

that if *m* ∪ = or ⊐ M ▼,

then will *m* ◆ =, or ⊐ M ■;


and therefore, by the fifth definition, we infer


that ∪ : ▼ : ◆ : ■.

∴ If four magnitudes, &c.



Harvard University - Houghton Library / pga_typ_805_47_3730


IF *the firſt be the ſame multiple of the ſecond, or the ſame part of it, that the third is of the fourth; the firſt is to the ſecond, as the third is to the fourth.*


Let  , the firſt, be the ſame multiple of  , the ſecond,

that  , the third, is of  , the fourth.

Then  :  ::  : 

take M  , *m*  , M  , *m*  ;

becauſe  is the ſame multiple of 

that  is of  (according to the hypotheſis);

and M  is taken the ſame multiple of 

that M  is of  ,

∴ (according to the third propoſition),


M  is the ſame multiple of

that M  is of  .

Harvard University - Houghton Library / pga_typ_805_47_3730

Therefore, if M  be of  a greater multiple than


m is, then M  is a greater multiple of  than

m  is; that is, if M  be greater than *m*  , then

M  will be greater than *m*  ; in the ſame manner

it can be ſhewn, if M  be equal *m*  , then

M  will be equal *m*  .

And, generally, if M  ⊏, = or ⊐ *m* 

then M  will be ⊏, = or ⊐ *m*  ;

∴ by the fifth definition,


 :  ::  :  .

Next, let  be the ſame part of 

that  is of  .

In this caſe alſo  :  ::  :  .

For, becauſe

 is the ſame part of  that  is of  ,

therefore  is the ſame multiple of 

that  is of  .

Therefore, by the preceding caſe,



and ∴  ,

by propoſition B.

∴ If the firſt be the ſame multiple, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *the firſt be to the ſecond as the third to the fourth, and if the firſt be a multiple, or a part of the ſecond; the third is the ſame multiple, or the ſame part of the fourth.*

Let  ; ■ ::  ;  ;

and firſt, let  be a multiple ■;

 ſhall be the ſame multiple of  .

First.	Second.	Third.	Fourth.
	■		

Take  =  .

Whatever multiple  is of ■

take  the ſame multiple of  ,


then, becauſe  : ■ ::  : 

and of the ſecond and fourth, we have taken equimultiples,

 and  , therefore (B. 5. pr. 4),

Harvard University - Houghton Library / pga_typ_805_47_3730


, but (conſt.),



and  is the ſame multiple of 

that  is of .

Next, let  :  ::  :  ,

and alſo  a part of  ;

then  ſhall be the ſame part of  .

Inverſely (B. 5.),  :  ::  :  ,

but  is a part of  ;

that is,  is a multiple of ;



∴ by the preceding caſe, is the ſame multiple of

that is,  is the ſame part of 

that  is of  .

∴ If the firſt be to the ſecond, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

EQUAL *magnitudes have the ſame ratio to the ſame magnitude, and the ſame has the ſame ratio to equal magnitudes.*

Let ● = ◆ and ■ any other magnitude;

then ● : ■ = ◆ : ■ and ■ : ● = ■ : ◆.

Becauſe ● = ◆,

∴ M ● = M ◆;

∴ if M ● ⊏, = or ⊐ *m* ■, then

M ◆ ⊏, = or ⊐ *m* ■,

and ∴ ● : ■ = ◆ : ■ (B. 5. def. 5).

From the foregoing reaſoning it is evident that,

if *m* ■ ⊏, = or ⊐ M ●, then

m ■ ⊏, = or ⊐ M ◆

∴ ■ : ● = ■ : ◆ (B. 5. def. 5).


∴ Equal magnitudes, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730


DEFINITION VII.

When of the equimultiples of four magnitudes (taken as in the fifth definition), the multiple of the firſt is greater than that of the ſecond, but the multiple of the third is not greater than the multiple of the fourth; then the firſt is ſaid to have to the ſecond a greater ratio than the third magnitude has to the fourth: and, on the contrary, the third is ſaid to have to the fourth a leſs ratio than the firſt has to the ſecond.

If, among the equimultiples of four magnitudes, compared as in the fifth definition, we ſhould find






, but



, or if we ſhould find any particular multiple M′ of the firſt and third, and a particular multiple *m′* of the ſecond and fourth, ſuch, that M′ times the firſt is ⊏ *m′* times the ſecond, but M′ times the third is not ⊏ *m′* times the fourth, *i. e.* = or ⊐ *m′* times the fourth; then the firſt is ſaid to have to the ſecond a greater ratio than the third has to the fourth; or the third has to the fourth, under ſuch circumſtances, a leſs ratio than the firſt has to the ſecond: although ſeveral other equimultiples may tend to ſhow that the four magnitudes are proportionals.

This definition will in future be expreſſed thus:—

If M′  *m′*  , but M′  or ⊐ *m′*  ,

then :  : .

In the above general expreſſion, M′ and *m′* are to be conſidered particular multiples, not like the multiples M

Harvard University - Houghton Library / pga_typ_805_47_3730

and *m* introduced in the fifth definition, which are in that definition conſidered to be every pair of multiples that can be taken. It muſt alſo be here obſerved, that ♥, ∪, ■, and the like ſymbols are to be conſidered merely the repreſentatives of geometrical magnitudes.

In a partial arithmetical way, this may be ſet forth as follows:

Let us take the four numbers, 8, 7, 10, and 9.

Firſt.	*Second.*	*Third.*	*Fourth.*
8	7	10	9
16	14	20	18
24	21	30	27
32	28	40	36
40	35	50	45
48	42	60	54
56	49	70	63
64	56	80	72
72	63	90	81
80	70	100	90
88	77	110	99
96	84	120	108
104	91	130	117
112	98	140	126
&c.	&c.	&c	&c.

Among the above multiples we find 16 ⊏ 14 and 20 ⊏ 18; that is, twice the firſt is greater than twice the ſecond, and twice the third is greater than twice the fourth; and 16 ⊐ 21 and 20 ⊐ 27; that is, twice the firſt is leſs than three times the ſecond, and twice the third is leſs than three times the fourth; and among the ſame multiples we can find 72 ⊏ 56 and 90 ⊏ 72: that is, 9 times the firſt is greater than 8 times the ſecond, and 9 times the third is greater than 8 times the fourth. Many other equimul-

Harvard University - Houghton Library / pga_typ_805_47_3730

tiples might be selected, which would tend to ſhow that the numbers 8, 7, 10, 9, were proportionals, but they are not, for we can find a multiple of the firſt ⊏ a multiple of the ſecond, but the ſame multiple of the third that has been taken of the firſt not ⊏ the ſame multiple of the fourth which has been taken of the ſecond; for inſtance, 9 times the firſt is ⊏ 10 times the ſecond, but 9 times the third is not ⊏ 10 times the fourth, that is, 72 ⊏ 70, but 90 not ⊏ 90, or 8 times the firſt we find ⊏ 9 times the ſecond, but 8 times the third is not greater than 9 times the fourth, that is, 64 ⊏ 63, but 80 is not ⊏ 81. When any ſuch multiples as theſe can be found, the firſt (8) is ſaid to have to the ſecond (7) a greater ratio than the third (10) has to the fourth (9), and on the contrary the third (10) is ſaid to have to the fourth (9) a leſs ratio than the firſt (8) has to the ſecond (7).

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *unequal magnitudes the greater has a greater ratio to the ſame than the leſs has: and the ſame magnitude has a greater ratio to the leſs than it has to the greater.*

Let ▲ and ■ be two unequal magnitudes,

and ● any other.

We ſhall firſt prove that ▲ which is the greater of the two unequal magnitudes, has a greater ratio to ● than ■, the leſs, has to ● ;

that is, ▲ : ● ⊏ ■ : ● ;

take M′ ▲, *m′* ●, M′ ■, and *m′* ● ;

ſuch, that M′ ▲ and M′ ■ ſhall be each ⊏ ● ;

alſo take *m′* ● the leaſt multiple of ●,

which will make *m′* ● ⊏ M′ ■ = M′ ▲ ;

∴ M′ ■ is not ⊏ *m′* ●,

but M′ ▲ is ⊏ *m′* ●, for,

as *m′* ● is the firſt multiple which firſt becomes ⊏ M′ ▲,

than (*m′* minus 1) ● or *m′* ● minus ● is not ⊏ M′ ▲,

and ● is not ⊏ M′ ■,

∴ *m′* ● minus ● + ● muſt be ⊐ M′ ▲ + M′ ■ ;

that is, *m′* ● muſt be ⊐ M′ ▲ ;

∴ M′ ▲ is ⊏ *m′* ● ; but it has been ſhown above that

M′ ■ is not ⊏ *m′* ● , therefore, by the ſeventh definition,

▲
■ has to ● a greater ratio than ■ : ● .

Next we ſhall prove that ● has a greater ratio to ■ , the

leſs, than it has to ▲■ , the greater ;

or, ● : ■ ⊏ ● : ▲■ .

Take *m′* ● , M′ ■ , *m′* ● , and M′ ▲■ ,
the ſame as in the firſt caſe, ſuch, that
M′ ▲ and M′ ■ will be each ⊏ ● , and *m′* ● the leaſt
multiple of ● , which firſt becomes greater
than M′ ■ = M′ ■ .

∴ *m′* ● minus ● is not ⊏ M′ ■ ,
and ● is not ⊏ M′ ▲ ; conſequently
m′ ● minus ● + ● is ⊐ M′ ■ + M′ ▲ ;

▲
∴ *m′* ● is ⊐ M′ ■ , and ∴ by the ſeventh definition,
▲
● has to ■ a greater ratio than ● has to ■ .

∴ Of unequal magnitudes, &c.

The contrivance employed in this propoſition for finding among the multiples taken, as in the fifth definition, a multiple of the firſt greater than the multiple of the ſecond, but the ſame multiple of the third which has been taken of the firſt, not greater than the ſame multiple of the fourth which has been taken of the ſecond, may be illuſtrated numerically as follows :—

The number 9 has a greater ratio to 7 than 8 has to 7 : that is, 9 : 7 ⊏ 8 : 7 ; or, 8 + 1 : 7 ⊏ 8 : 7 .

Harvard University - Houghton Library / pga_typ_805_47_3730

The multiple of 1, which firſt becomes greater than 7, is 8 times, therefore we may multiply the firſt and third by 8, 9, 10, or any other greater number; in this caſe, let us multiply the firſt and third by 8, and we have 64 + 8 and 64: again, the firſt multiple of 7 which becomes greater than 64 is 10 times; then, by multiplying the ſecond and fourth by 10, we ſhall have 70 and 70; then, arranging theſe multiples, we have—

8 times the first.	10 times the second.	8 times the third.	10 times the fourth.
64 + 8	70	64	70

Conſequently 64 + 8, or 72, is greater than 70, but 64 is not greater than 70, ∴ by the ſeventh definition, 9 has a greater ratio to 7 than 8 has to 7.

The above is merely illuſtrative of the foregoing demonſtration, for this property could be ſhown of theſe or other numbers very readily in the following manner; becauſe, if an antecedent contains its conſequent a greater number of times than another antecedent contains its conſequent, or when a fraction is formed of an antecedent for the numerator, and its conſequent for the denominator be greater than another fraction which is formed of another antecedent for the numerator and its conſequent for the denominator, the ratio of the firſt antecedent to its conſequent is greater than the ratio of the laſt antecedent to its conſequent.

Thus, the number 9 has a greater ratio to 7, than 8 has to 7, for $\frac{9}{7}$ is greater than $\frac{8}{7}$.

Again, 17 : 19 is a greater ratio than 13 : 15, becauſe $\frac{17}{19} = \frac{17 \times 15}{19 \times 15} = \frac{255}{285}$, and $\frac{13}{15} = \frac{13 \times 19}{15 \times 19} = \frac{247}{285}$, hence it is evident that $\frac{255}{285}$ is greater than $\frac{247}{285}$, ∴ $\frac{17}{19}$ is greater than

Harvard University - Houghton Library / pga_typ_805_47_3730

$\frac{13}{15}$, and, according to what has been above ſhown, 17 has to 19 a greater ratio than 13 has to 15.

So that the general terms upon which a greater, equal, or leſs ratio exiſts are as follows:—

If $\frac{A}{B}$ be greater than $\frac{C}{D}$, A is ſaid to have to B a greater ratio than C has to D; if $\frac{A}{B}$ be equal to $\frac{C}{D}$, then A has to B the ſame ratio which C has to D; and if $\frac{A}{B}$ be leſs than $\frac{C}{D}$, A is ſaid to have to B a leſs ratio than C has to D.

The ſtudent ſhould underſtand all up to this propoſition perfectly before proceeding further, in order fully to comprehend the following propoſitions of this book. We therefore ſtrongly recommend the learner to commence again, and read up to this ſlowly, and carefully reaſon at each ſtep, as he proceeds, particularly guarding againſt the miſchievous ſyſtem of depending wholly on the memory. By following theſe inſtructions, he will find that the parts which uſually preſent conſiderable difficulties will preſent no difficulties whatever, in proſecuting the ſtudy of this important book.

Harvard University - Houghton Library / pga_typ_805_47_3730

MAGNITUDES *which have the ſame ratio to the ſame magnitude are equal to one another; and thoſe to which the ſame magnitude has the ſame ratio are equal to one another.*

Let ◆ : ■ : : ● : ■, then ◆ = ●.

For, if not, let ◆ ⊏ ●, then will

◆ : ■ ⊏ ● : ■ (B. 5. pr. 8),

which is abſurd according to the hypotheſis.

∴ ◆ is not ⊏ ●.

In the ſame manner it may be ſhown, that

● is not ⊏ ◆,

∴ ◆ = ●.

Again, let ■ : ◆ :: ■ : ●, then will ◆ = ●.

For (invert.) ◆ : ■ :: ● : ■,

therefore, by the firſt caſe, ◆ = ●.

∴ Magnitudes which have the ſame ratio, &c.

This may be ſhown otherwiſe, as follows :—

Let A : B = A : C, then B = C, for, as the fraction $\frac{A}{B}$ = the fraction $\frac{A}{C}$, and the numerator of one equal to the numerator of the other, therefore the denominator of theſe fractions are equal, that is B = C.

Again, if B : A = C : A, B = C. For, as $\frac{B}{A} = \frac{C}{A}$, B muſt = C.

Harvard University - Houghton Library / pga_typ_805_47_3730

THAT *magnitude which has a greater ratio than another has unto the ſame magnitude, is the greater of the two: and that magnitude to which the ſame has a greater ratio than it has unto another magnitude, is the leſs of the two.*

Let ▼ : ■ ⊏ ● : ■, then ▼ ⊏ ●.

For if not, let ▼ = or ⊐ ● ;

then, ▼ : ■ = ● : ■ (B. 5. pr. 7) or

▼ : ■ ⊐ ● : ■ (B. 5. pr. 8) and (invert.),

which is abſurd according to the hypotheſis.

∴ ▼ is not = or ⊐ ●, and

∴ ▼ muſt be ⊏ ●.

Again, let ■ : ● ⊏ ■ : ▼,

then, ● ⊐ ▼.

For if not, ● muſt be ⊏ or = ▼,

then ■ : ● ⊐ ■ : ▼ (B. 5. pr. 8) and (invert.);

or ■ : ● = ■ : ▼ (B. 5. pr. 7), which is abſurd (hyp.);

∴ ● is not ⊏ or = ▼,

and ∴ ● muſt be ⊐ ▼.

∴ That magnitude which has, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

RATIOS *that are the ſame to the ſame ratio, are the ſame to each other.*

Let ◆ : ■ = ● : ⬟ and ● : ⬟ = ▲ : ●,

then will ◆ : ■ = ▲ : ●.

For if M ◆ ⊏, =, or ⊐ *m* ■,

then M ● ⊏, =, or ⊐ *m* ⬟,

and if M ● ⊏, =, or ⊐ *m* ⬟,

then M ▲ ⊏, =, or ⊐ *m* ●, (B. 5. def. 5);

∴ if M ◆ ⊏, =, or ⊐ *m* ■, M ▲ ⊏, =, or ⊐ *m* ●,

and ∴ (B. 5. def. 5) ◆ : ■ = ▲ : ●.

∴ Ratios that are the ſame, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *any number of magnitudes be proportionals, as one of the antecedents is to its conſequent, ſo ſhall all the antecedents taken together be to all the conſequents.*

Let ■ : ● = ⋃ : ◇ = ◆ : ⬟ = • : ▾ = ▲ : ●;

then will ■ : ● =

■ + ⋃ + ◆ + • + ▲ : ● + ◇ + ⬟ + ▾ + ●.

For if M ■ ⊏ *m* ●, then M ⋃ ⊏ *m* ◇,

and M ◆ ⊏ *m* ⬟ M • ⊏ *m* ▾,

alſo M ▲ ⊏ *m* ●. (B. 5. def. 5.)

Therefore, if M ■ ⊏ *m* ●, then will

M ■ + M ⋃ + M ◆ + M • + M ▲,

or M (■ + ⋃ + ◆ + • + ▲) be greater

than *m* ● + *m* ◇ + *m* ⬟ + *m* ▾ + *m* ●,

or *m* (● + ◇ + ⬟ + ▾ + ●).

In the ſame way it may be ſhown, if M times one of the antecedents be equal to or leſs than *m* times one of the conſequents, M times all the antecedents taken together, will be equal to or leſs than *m* times all the conſequents taken together. Therefore, by the fifth definition, as one of the antecedents is to its conſequent, ſo are all the antecedents taken together to all the conſequents taken together.

∴ If any number of magnitudes, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *the firſt has to the ſecond the ſame ratio which the third has to the fourth, but the third to the fourth a greater ratio than the fifth has to the ſixth; the firſt ſhall alſo have to the ſecond a greater ratio than the fifth to the ſixth.*

Let ▼ : ∪ = ■ : ◆, but ■ : ◆ ⊏ ◇ : ●,

then ▼ : ∪ ⊏ ◇ : ●.

For, becauſe ■ : ◆ ⊏ ◇ : ●, there are ſome multiples (M′ and *m*′) of ■ and ◇, and of ◆ and ●, ſuch that M′ ■ ⊏ *m*′ ◆,

but M′ ◇ not ⊏ *m*′ ●, by the ſeventh definition.

Let theſe multiples be taken, and take the ſame multiples of ▼ and ∪.

∴ (B. 5. def. 5.) if M′ ▼ ⊏, =, or ⊐ *m*′ ∪;

then will M′ ■ ⊏, =, or ⊐ *m*′ ◆,

but M′ ■ ⊏ *m*′ ◆ (conſtruction);

∴ M′ ▼ ⊏ *m*′ ∪,

but M′ ◇ is not ⊏ *m*′ ● (conſtruction);

and therefore by the ſeventh definition,

▼ : ∪ ⊏ ◇ : ●.

∴ If the firſt has to the ſecond, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *the firſt has the ſame ratio to the ſecond which the third has to the fourth; then, if the firſt be greater than the third, the ſecond ſhall be greater than the fourth; and if equal, equal; and if leſs, leſs.*

Let ▼ : ∪ :: ■ : ◆, and firſt ſuppoſe

▼ ⊏ ■, then will ∪ ⊏ ◆.

For ▼ : ∪ ⊏ ■ : ∪ (B. 5. pr. 8), and by the

hypotheſis, ▼ : ∪ = ■ : ◆ ;

∴ ■ : ◆ ⊏ ■ : ∪ (B. 5. pr. 13),

∴ ◆ ⊐ ∪ (B. 5. pr. 10.), or ∪ ⊏ ◆.

Secondly, let ▼ = ■, then will ∪ = ◆.

For ▼ : ∪ = ■ : ∪ (B. 5. pr. 7),

and ▼ : ∪ = ■ : ◆ (hyp.);

∴ ■ : ∪ = ■ : ◆ (B. 5. pr. 11),

and ∴ ∪ = ◆ (B. 5, pr. 9).

Thirdly, if ▼ ⊐ ■, then will ∪ ⊐ ◆ ;

becauſe ■ ⊏ ▼ and ■ : ◆ = ▼ : ∪ ;

∴ ◆ ⊏ ∪, by the firſt caſe,

that is, ∪ ⊐ ◆.

∴ If the firſt has the ſame ratio, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

MAGNITUDES *have the ſame ratio to one another which their equimultiples have.*

Let ● and ■ be two magnitudes;

then, ● : ■ :: M′ ● : M′ ■.

For ● : ■ = ● : ■

= ● : ■

= ● : ■

∴ ● : ■ :: 4 ● : 4 ■. (B. 5. pr. 12).

And as the ſame reaſoning is generally applicable, we have

● : ■ :: M′ ● : M′ ■.

∴ Magnitudes have the ſame ratio, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

DEFINITION XIII.

THE technical term permutando, or alternando, by permutation or alternately, is uſed when there are four proportionals, and it is inferred that the firſt has the ſame ratio to the third which the ſecond has to the fourth; or that the firſt is to the third as the ſecond is to the fourth: as is ſhown in the following propoſition:—

Let ● : ◆ :: ▼ : ■,

by "permutando" or "alternando" it is

inferred ● : ▼ :: ◆ : ■.

It may be neceſſary here to remark that the magnitudes ●, ◆, ▼, ■, muſt be homogeneous, that is, of the ſame nature or ſimilitude of kind; we muſt therefore, in ſuch caſes, compare lines with lines, ſurfaces with ſurfaces, ſolids with ſolids, &c. Hence the ſtudent will readily perceive that a line and a ſurface, a ſurface and a ſolid, or other heterogenous magnitudes, can never ſtand in the relation of antecedent and conſequent.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *four magnitudes of the ſame kind be proportionals, they are alſo proportionals when taken alternately.*

Let ▼ : ▭ :: ■ : ◆, then ▼ : ■ :: ▭ : ◆.

For M ▼ : M ▭ :: ▼ : ▭ (B. 5. pr. 15),

and M ▼ : M ▭ :: ■ : ◆ (hyp.) and (B. 5. pr. 11);

alſo *m* ■ : *m* ◆ :: ■ : ◆ (B. 5. pr. 15);

∴ M ▼ : M ▭ :: *m* ■ : *m* ◆ (B. 5. pr. 14),

and ∴ if M ▼ ⊏, =, or ⊐ *m* ■,

then will M ▭ ⊏, =, or ⊐ *m* ◆ (B. 5. pr. 14);

therefore, by the fifth definition,

▼ : ■ :: ▭ : ◆.

∴ If four magnitudes of the ſame kind, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

DEFINITION XVI.

DIVIDENDO, by diviſion, when there are four proportionals, and it is inferred, that the exceſs of the firſt above the ſecond is to the ſecond, as the exceſs of the third above the fourth, is to the fourth.

Let A : B :: C : D;

by "dividendo" it is inferred

A minus B : B :: C minus D : D.

According to the above, A is ſuppoſed to be greater than B, and C greater than D; if this be not the caſe, but to have B greater than A, and D greater than C, B and D can be made to ſtand as antecedents, and A and C as conſequents, by "invertion"

B : A :: D : C;

then, by "dividendo," we infer

B minus A : A :: D minus C : C.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *magnitudes, taken jointly, be proportionals, they ſhall alſo be proportionals when taken ſeparately: that is, if two magnitudes together have to one of them the ſame ratio which two others have to one of theſe, the remaining one of the firſt two ſhall have to the other the ſame ratio which the remaining one of the laſt two has to the other of theſe.*

Let ♥ + ⋃ : ⋃ :: ■ + ◆ : ◆,

then will ♥ : ⋃ :: ■ : ◆.

Take M ♥ ⊏ *m* ⋃ to each add M ⋃,

then we have M ♥ + M ⋃ ⊏ *m* ⋃ + M ⋃,

or M (♥ + ⋃) ⊏ (*m* + M) ⋃:

but becauſe ♥ + ⋃ : ⋃ :: ■ + ◆ : ◆ (hyp.),

and M (♥ + ⋃) ⊏ (*m* + M) ⋃;

∴ M (■ + ◆) ⊏ (*m* + M) ◆ (B. 5. def. 5);

∴ M ■ + M ◆ ⊏ *m* ◆ + M ◆;

∴ M ■ ⊏ *m* ◆, by taking M ◆ from both ſides:

that is, when M ♥ ⊏ *m* ⋃, then M ■ ⊏ *m* ◆.

In the ſame manner it may be proved, that if

M ♥ = or ⊐ *m* ⋃, then will M ■ = or ⊐ *m* ◆;

and ∴ ♥ : ⋃ : : ■ : ◆ (B. 5. def. 5).

∴ If magnitudes taken jointly, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

DEFINITION XV.

THE term componendo, by compoſition, is uſed when there are four proportionals; and it is inferred that the firſt together with the ſecond is to the ſecond as the third together with the fourth is to the fourth.

Let A : B :: C : D;

then, by the term "componendo," it is inferred that

$$A + B : B :: C + D : D.$$

By "invertion" B and D may become the firſt and third, A and C the ſecond and fourth, as

B : A :: D : C,

then, by "componendo," we infer that

$$B + A : A :: D + C : C.$$

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *magnitudes, taken ſeparately, be proportionals, they ſhall alſo be proportionals when taken jointly: that is, if the firſt be to the ſecond as the third is to the fourth, the firſt and ſecond together ſhall be to the ſecond as the third and fourth together is to the fourth.*

Let ▼ : ⋃ :: ■ : ◆,

then ▼ + ⋃ : ⋃ :: ■ + ◆ : ◆ ;

for if not, let ▼ + ⋃ : ⋃ :: ■ + ● : ●,

ſuppoſing ● not = ◆ ;

∴ ▼ : ⋃ :: ■ : ● (B. 5. pr. 17);

but ▼ : ⋃ :: ■ : ◆ (hyp.);

∴ ■ : ● :: ■ : ◆ (B. 5. pr. 11);

∴ ● = ◆ (B. 5. pr. 9),

which is contrary to the ſuppoſition;

∴ ● is not unequal to ◆ ;

that is ● = ◆ ;

∴ ▼ + ⋃ : ⋃ :: ■ + ◆ : ◆.

∴ If magnitudes, taken ſeparately, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *a whole magnitude be to a whole, as a magnitude taken from the firſt, is to a magnitude taken from the other; the remainder ſhall be to the remainder, as the whole to the whole.*

Let ♥ + ∪ : ■ + ◆ :: ♥ : ■,

then will ∪ : ◆ :: ♥ + ∪ : ■ + ◆,

For ♥ + ∪ : ♥ :: ■ + ◆ : ■ (alter.),

∴ ∪ : ♥ :: ◆ : ■ (divid.),

again ∪ : ◆ :: ♥ : ■ (alter.),

but ♥ + ∪ : ■ + ◆ :: ♥ : ■ hyp.);

therefore ∪ : ◆ :: ♥ + ∪ : ■ + ◆

(B. 5. pr. 11).

∴ If a whole magnitude be to a whole, &c.

DEFINITION XVII.

THE term "convertendo," by converſion, is made uſe of by geometricians, when there are four proportionals, and it is inferred, that the firſt is to its exceſs above the ſecond, as the third is to its exceſs above the fourth. See the following propoſition:—

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *four magnitudes be proportionals, they are alſo proportionals by converſion: that is, the firſt is to its exceſs above the ſecond, as the third to its exceſs above the fourth.*

Let ● ◇ : ◇ :: ■ ◆ : ◆,

then ſhall ● ◇ : ● :: ■ ◆ : ■,

Becauſe ● ◇ : ◇ : ■ ◆ : ◆;

therefore ● : ◇ :: ■ : ◆ (divid.),

∴ ◇ : ● :: ◆ : ■ (inver.),

∴ ● ◇ : ● :: ■ ◆ : ■ (compo.).

∴ If four magnitudes, &c.

DEFINITION XVIII.

"Ex æquali" (ſc. diſtantiâ), or ex æquo, from equality of diſtance: when there is any number of magnitudes more than two, and as many others, ſuch that they are proportionals when taken two and two of each rank, and it is inferred that the firſt is to the laſt of the firſt rank of magnitudes, as the firſt is to the laſt of the others: "of this there are the two following kinds, which ariſe from the different order in which the magnitudes are taken, two and two."

Harvard University - Houghton Library / pga_typ_805_47_3730

DEFINITION XIX.

" Ex æquali," from equality. This term is uſed ſimply by itſelf, when the firſt magnitude is to the ſecond of the firſt rank, as the firſt to the ſecond of the other rank; and as the ſecond is to the third of the firſt rank, ſo is the ſecond to the third of the other; and ſo on in order: and the inference is as mentioned in the preceding definition; whence this is called ordinate proportion. It is demonſtrated in Book 5. pr. 22.

Thus, if there be two ranks of magnitudes,

A, B, C, D, E, F, the firſt rank,

and L, M, N, O, P, Q, the ſecond,

ſuch that A : B :: L : M, B : C :: M : N,

C : D :: N : O, D : E :: O : P, E : F :: P : Q;

we infer by the term " ex æquali" that

A : F :: L : Q.

Harvard University - Houghton Library / pga_typ_805_47_3730

DEFINITION XX.

"Ex æquali in proportione perturbatâ ſeu inordinatâ," from equality in perturbate, or diſorderly proportion. This term is uſed when the firſt magnitude is to the ſecond of the firſt rank as the laſt but one is to the laſt of the ſecond rank; and as the ſecond is to the third of the firſt rank, ſo is the laſt but two to the laſt but one of the ſecond rank; and as the third is to the fourth of the firſt rank, ſo is the third from the laſt to the laſt but two of the ſecond rank; and ſo on in a croſs order: and the inference is in the 18th definition. It is demonſtrated in B. 5. pr. 23.

Thus, if there be two ranks of magnitudes,

A, B, C, D, E, F, the firſt rank,

and L, M, N, O, P, Q, the ſecond,

ſuch that A : B :: P : Q, B : C :: O : P,

C : D :: N : O, D : E :: M : N, E : F :: L : M;

the term "ex æquali in proportione perturbatâ ſeu inordinatâ" infers that

A : F :: L : Q.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *there be three magnitudes, and other three, which, taken two and two, have the ſame ratio; then, if the firſt be greater than the third, the fourth ſhall be greater than the ſixth; and if equal, equal; and if leſs, leſs.*

Let ▼, ◡, ■, be the firſt three magnitudes,

and ◆, ◇, ●, be the other three,

ſuch that ▼ : ◡ :: ◆ : ◇, and ◡ : ■ :: ◇ : ●.

Then, if ▼ ⊏, =, or ⊐ ■, then will ◆ ⊏, =,

or ⊐ ●.

From the hypotheſis, by alternando, we have

▼ : ◆ :: ◡ : ◇,

and ◡ : ◇ :: ■ : ●;

∴ ▼ : ◆ :: ■ : ● (B. 5. pr. 11);

∴ if ▼ ⊏, =, or ⊐ ■, then will ◆ ⊏, =,

or ⊐ ● (B. 5. pr. 14).

∴ If there be three magnitudes, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *there be three magnitudes, and other three which have the ſame ratio, taken two and two, but in a croſs order; then if the firſt magnitude be greater than the third, the fourth ſhall be greater than the ſixth; and if equal, equal; and if leſs, leſs.*

Let ▼, ▲, ■, be the firſt three magnitudes,

and ◆, ◇, ●, the other three,

ſuch that ▼ : ▲ :: ◇ : ●, and ▲ : ■ :: ◆ : ◇.

Then, if ▼ ⊏, =, or ⊐ ■, then

will ◆ ⊏, =, ⊐ ●.

Firſt, let ▼ be ⊏ ■ :

then, becauſe ▲ is any other magnitude,

▼ : ▲ ⊏ ■ : ▲ (B. 5. pr. 8);

but ◇ : ● :: ▼ : ▲ (hyp.);

∴ ◇ : ● ⊏ ■ : ▲ (B. 5. pr. 13);

and becauſe ▲ : ■ :: ◆ : ◇ (hyp.);

∴ ■ : ▲ :: ◇ : ◆ (inv.),

and it was ſhown that ◇ : ● ⊏ ■ : ▲,

∴ ◇ : ● ⊏ ◇ : ◆ (B. 5. pr. 13);

Harvard University - Houghton Library / pga_typ_805_47_3730

$\therefore$ ● ⊐ ◆,

that is ◆ ⊏ ●.

Secondly, let ▼ = ■; then ſhall ◆ = ●.

For becauſe ▼ = ■,

▼ : ▲ = ■ : ▲ (B. 5. pr. 7);

but ▼ : ▲ = ◇ : ● (hyp.),

and ■ : ▲ = ◇ : ◆ (hyp. and inv.),

$\therefore$ ◇ : ● = ◇ : ◆ (B. 5. pr. 11),

$\therefore$ ◆ = ● (B. 5. pr. 9).

Next, let ▼ be ⊐ ■, then ◆ ſhall be ⊐ ●;

for ■ ⊏ ▼,

and it has been ſhown that ■ : ▲ = ◇ : ◆,

and ▲ : ▼ = ● : ◇;

$\therefore$ by the firſt caſe ● is ⊏ ◆,

that is, ◆ ⊐ ●.

$\therefore$ If there be three, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *there be any number of magnitudes, and as many others, which, taken two and two in order, have the ſame ratio; the firſt ſhall have to the laſt of the firſt magnitudes the ſame ratio which the firſt of the others has to the laſt of the ſame.*

N.B.—*This is uſually cited by the words "ex æquali," or "ex æquo."*

Firſt, let there be magnitudes ▼, ◆, ■,

and as many others ◆, ◇, ●,

ſuch that

▼ : ◆ :: ◆ : ◇,

and ◆ : ■ :: ◇ : ● ;

then ſhall ▼ : ■ :: ◆ : ●.

Let theſe magnitudes, as well as any equimultiples whatever of the antecedents and conſequents of the ratios, ſtand as follows:—

▼, ◆, ■, ◆, ◇, ●,

and

M ▼, *m* ◆, N ■, M ◆, *m* ◇, N ●,

becauſe ▼ : ◆ :: ◆ : ◇ ;

∴ M ▼ : *m* ◆ :: M ◆ : *m* ◇ (B. 5. p. 4).

For the ſame reaſon

m ◆ : N ■ :: *m* ◇ : N ● ;

and becauſe there are three magnitudes,

Harvard University - Houghton Library / pga_typ_805_47_3730

M ▼, *m* ◆, N ■,

and other three, M ◆, *m* ◇, N ●,

which, taken two and two, have the ſame ratio;

∴ if M ▼ ⊏, =, or ⊐ N ■

then will M ◆ ⊏, =, or ⊐ N ●, by (B. 5. pr. 20);

and ∴ ▼ : ■ :: ◆ : ● (def. 5).

Next, let there be four magnitudes, ⬟, ◆, ■, ◆,

and other four, ◇, ●, ▬, ▲,

which, taken two and two, have the ſame ratio,

that is to ſay, ⬟ : ◆ :: ◇ : ●,

◆ : ■ :: ● : ▬,

and ■ : ◆ :: ▬ : ▲,

then ſhall ⬟ : ◆ :: ◇ : ▲;

for, becauſe ⬟, ◆, ■, are three magnitudes,

and ◇, ●, ▬, other three,

which, taken two and two, have the ſame ratio;

therefore, by the foregoing caſe, ⬟ : ■ :: ◇ : ▬,

but ■ : ◆ :: ▬ : ▲;

therefore again, by the firſt caſe, ⬟ : ◆ :: ◇ : ▲;

and ſo on, whatever the number of magnitudes be.

∴ If there be any number, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *there be any number of magnitudes, and as many others, which, taken two and two in a croſs order, have the ſame ratio; the firſt ſhall have to the laſt of the firſt magnitudes the ſame ratio which the firſt of the others has to the laſt of the ſame.*

N.B.—*This is uſually cited by the words "ex æquali in proportione perturbatâ;" or "ex æquo perturbato."*

Firſt, let there be three magnitudes, ▼, ▢, ■,

and other three, ◆, ◇, ●,

which, taken two and two in a croſs order,

have the ſame ratio;

that is, ▼ : ▢ :: ◇ : ●,

and ▢ : ■ :: ◆ : ◇,

then ſhall ▼ : ■ :: ◆ : ●.

Let theſe magnitudes and their reſpective equimultiples be arranged as follows:—

▼, ▢, ■, ◆, ◇, ●,

M▼, M▢, *m*■, M◆, *m*◇, *m*●,

then ▼ : ▢ :: M▼ : M▢ (B. 5. pr. 15);

and for the ſame reaſon

◇ : ● :: *m*◇ : *m*● ;

but ▼ : ▢ :: ◇ : ● (hyp.),

Harvard University - Houghton Library / pga_typ_805_47_3730

∴ M ▼ : M ▽ :: ◇ : ● (B. 5. pr. 11);

and becauſe ▽ : ■ :: ◆ : ◇ (hyp.),

∴ M ▽ : *m* ■ :: ◆ : *m* ◇ (B. 5. pr. 4);

then, becauſe there are three magnitudes,

M ▼, M ▽, *m* ■,

and other three, M ◆, *m* ◇, *m* ●,

which, taken two and two in a croſs order, have the ſame ratio;

therefore, if M ▼ ⊏, =, or ⊐ *m* ■,

then will M ◆ ⊏, =, or ⊐ *m* ● (B. 5. pr. 21),

and ∴ ▼ : ■ :: ◆ : ● (B. 5. def. 5).

Next, let there be four magnitudes,

▼, ▽, ■, ◆,

and other four, ◇, ●, ▬, ▲,

which, when taken two and two in a croſs order, have the ſame ratio; namely,

▼ : ▽ :: ▬ : ▲,

▽ : ■ :: ● : ▬,

and ■ : ◆ :: ◇ : ●,

then ſhall ▼ : ◆ :: ◇ : ▲.

For, becauſe ▼, ▽, ■ are three magnitudes,

Harvard University - Houghton Library / pga_typ_805_47_3730

and ●, ▬, ▲, other three,

which, taken two and two in a croſs order, have the ſame ratio,

therefore, by the firſt caſe, ▼ : ■ :: ● : ▲,


but ■ : ◆ :: ◇ : ●,

therefore again, by the firſt caſe, ▼ : ◆ :: ◇ : ▲;

and ſo on, whatever be the number of ſuch magnitudes.

∴ If there be any number, &c.

IF *the firſt has to the ſecond the ſame ratio which the third has to the fourth, and the fifth to the ſecond the ſame which the ſixth has to the fourth, the firſt and fifth together ſhall have to the ſecond the ſame ratio which the third and ſixth together have to the fourth.*



Let ▼ : ∪ :: ■ : ◆,

and ◇ : ∪ :: ● : ◆,

then ▼ + ◇ : ∪ :: ■ + ● : ◆.

For ◇ : ∪ :: ● : ◆ (hyp.),

and ∪ : ▼ :: ◆ : ■ (hyp.) and (invert.),

∴ ◇ : ▼ :: ● : ■ (B. 5. pr. 22);

and, becauſe theſe magnitudes are proportionals, they are proportionals when taken jointly,

∴ ▼ + ◇ : ◇ :: ● + ■ : ● (B. 5. pr. 18),

but ◇ : ∪ :: ● : ◆ (hyp.),

∴ ▼ + ◇ : ∪ :: ● + ■ : ◆ (B. 5. pr. 22).

∴ If the firſt, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *four magnitudes of the ſame kind are proportionals, the greateſt and leaſt of them together are greater than the other two together.*

Let four magnitudes, ♥ + ⊍, ■ + ◆, ⊍, and ◆,

of the ſame kind, be proportionals, that is to ſay,

♥ + ⊍ : ■ + ◆ :: ⊍ : ◆,

and let ♥ + ⊍ be the greateſt of the four, and conſequently by pr. A and 14 of Book 5, ◆ is the leaſt;

then will ♥ + ⊍ + ◆ be ⊏ ■ + ◆ + ⊍;

becauſe ♥ + ⊍ : ■ + ◆ :: ⊍ : ◆,

∴ ♥ : ■ :: ♥ + ⊍ : ■ + ◆ (B. 5. pr. 19),

but ♥ + ⊍ ⊏ ■ + ◆ (hyp.),

∴ ♥ ⊏ ■ (B. 5. pr. A);

to each of theſe add ⊍ + ◆,

∴ ♥ + ⊍ + ◆ ⊏ ■ + ⊍ + ◆.

∴ If four magnitudes, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

DEFINITION X.

When three magnitudes are proportionals, the firſt is ſaid to have to the third the duplicate ratio of that which it has to the ſecond.

For example, if A, B, C, be continued proportionals, that is, A : B :: B : C, A is ſaid to have to C the duplicate ratio of A : B ;

or $\frac{A}{C}$ = the ſquare of $\frac{A}{B}$.

This property will be more readily ſeen of the quantities

ar^2, ar, a, for $ar^2 : ar :: ar : a$;

and $\frac{ar^2}{a} = r^2$ = the ſquare of $\frac{ar^2}{ar} = r$,

or of a, ar, ar^2 ;

for $\frac{a}{ar^2} = \frac{1}{r^2}$ = the ſquare of $\frac{a}{ar} = \frac{1}{r}$.

DEFINITION XI.

When four magnitudes are continual proportionals, the firſt is ſaid to have to the fourth the triplicate ratio of that which it has to the ſecond; and ſo on, quadruplicate, &c. increaſing the denomination ſtill by unity, in any number of proportionals.

For example, let A, B, C, D, be four continued proportionals, that is, A : B :: B : C :: C : D; A is ſaid to have to D, the triplicate ratio of A to B ;

or $\frac{A}{D}$ = the cube of $\frac{A}{B}$.

Harvard University - Houghton Library / pga_typ_805_47_3730

This definition will be better underſtood, and applied to a greater number of magnitudes than four that are continued proportionals, as follows:—

Let ar^3, ar^2, ar, a, be four magnitudes in continued proportion, that is, $ar^3 : ar^2 :: ar^2 : ar :: ar : a$,

then $\frac{ar^3}{a} = r^3 =$ the cube of $\frac{ar^3}{ar^2} = r$.

Or, let ar^5, ar^4, ar^3, ar^2, ar, a, be ſix magnitudes in proportion, that is

$ar^5 : ar^4 :: ar^4 \cdot ar^3 :: ar^3 : ar^2 :: ar^2 : ar :: ar : a$,

then the ratio $\frac{ar^5}{a} = r^5 =$ the fifth power of $\frac{ar^5}{ar^4} = r$.

Or, let a, ar, ar^2, ar^3, ar^4, be five magnitudes in continued proportion; then $\frac{a}{ar^4} = \frac{1}{r^4} =$ the fourth power of $\frac{a}{ar} = \frac{1}{r}$.

DEFINITION A.

To know a compound ratio:—

When there are any number of magnitudes of the ſame kind, the firſt is ſaid to have to the laſt of them the ratio compounded of the ratio which the firſt has to the ſecond, and of the ratio which the ſecond has to the third, and of the ratio which the third has to the fourth; and ſo on, unto the laſt magnitude.

For example, if A, B, C, D, be four magnitudes of the ſame kind, the firſt A is ſaid to have to the laſt D the ratio compounded of the ratio of A to B, and of the ratio of B to C, and of the ratio of C to D; or, the ratio of



A to D is ſaid to be compounded of the ratios of A to B, B to C, and C to D.

And if A has to B the ſame ratio which E has to F, and B to C the ſame ratio that G has to H, and C to D the ſame that K has to L; then by this definition, A is said to have to D the ratio compounded of ratios which are the ſame with the ratios of E to F, G to H, and K to L. And the ſame thing is to be underſtood when it is more briefly expreſſed by ſaying, A has to D the ratio compounded of the ratios of E to F, G to H, and K to L.

In like manner, the ſame things being ſuppoſed; if M has to N the ſame ratio which A has to D, then for ſhortneſs ſake, M is ſaid to have to N the ratio compounded of the ratios of E to F, G to H, and K to L.

This definition may be better underſtood from an arithmetical or algebraical illuſtration; for, in fact, a ratio compounded of ſeveral other ratios, is nothing more than a ratio which has for its antecedent the continued product of all the antecedents of the ratios compounded, and for its conſequent the continued product of all the conſequents of the ratios compounded.

Thus, the ratio compounded of the ratios of

$$2 : 3,\ 4 : 7,\ 6 : 11,\ 2 : 5,$$

is the ratio of $2 \times 4 \times 6 \times 2 : 3 \times 7 \times 11 \times 5$,

or the ratio of $96 : 1155$, or $32 : 385$.

And of the magnitudes A, B, C, D, E, F, of the ſame kind, A : F is the ratio compounded of the ratios of

$$A : B,\ B : C,\ C : D,\ D : E,\ E : F;$$

for $A \times B \times C \times D \times E : B \times C \times D \times E \times F$,

or $\frac{A \times B \times C \times D \times E}{B \times C \times D \times E \times F} = \frac{A}{F}$, or the ratio of A : F.

Harvard University - Houghton Library / pga_typ_805_47_3730

RATIOS *which are compounded of the ſame ratios are the ſame to one another.*

Let A : B :: F : G,
B : C :: G : H,
C : D :: H : K,
and D : E :: K : L.



Then the ratio which is compounded of the ratios of A : B, B : C, C : D, D : E, or the ratio of A : E, is the ſame as the ratio compounded of the ratios of F : G, G : H, H : K, K : L, or the ratio of F : L.

$$\text{For } \frac{A}{B} = \frac{F}{G},$$
$$\frac{B}{C} = \frac{G}{H},$$
$$\frac{C}{D} = \frac{H}{K},$$
$$\text{and } \frac{D}{E} = \frac{K}{L};$$
$$\therefore \frac{A \times B \times C \times D}{B \times C \times D \times E} = \frac{F \times G \times H \times K}{G \times H \times K \times L},$$
$$\text{and } \therefore \frac{A}{E} = \frac{F}{L},$$

or the ratio of A : E is the ſame as the ratio of F : L.

The ſame may be demonſtrated of any number of ratios ſo circumſtanced.

Next, let A : B :: K : L,
B : C :: H : K,
C : D :: G : H,
D : E :: F : G.

Harvard University - Houghton Library / pga_typ_805_47_3730

Then the ratio which is compounded of the ratios of A : B, B : C, C : D, D : E, or the ratio of A : E, is the ſame as the ratio compounded of the ratios of K : L, H : K, G : H, F : G, or the ratio of F : L.

For $\frac{A}{B} = \frac{K}{L}$,

$\frac{B}{C} = \frac{H}{K}$,

$\frac{C}{D} = \frac{G}{H}$,

and $\frac{D}{E} = \frac{F}{G}$;

$\therefore \frac{A \times B \times C \times D}{B \times C \times D \times E} = \frac{K \times H \times G \times F}{L \times K \times H \times G}$,

and $\therefore \frac{A}{E} = \frac{F}{L}$,

or the ratio of A : E is the ſame as the ratio of F : L.

$\therefore$ Ratios which are compounded, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *ſeveral ratios be the ſame to ſeveral ratios, each to each, the ratio which is compounded of ratios which are the ſame to the firſt ratios, each to each, ſhall be the ſame to the ratio compounded of ratios which are the ſame to the other ratios, each to each.*

A B C D E F G H		P Q R S T
a b c d e f g h		V W X Y Z

If A : B :: *a* : *b*	and A : B :: P : Q	*a* : *b* :: V : W
C : D :: *c* : *d*	C : D :: Q : R	*c* : *d* :: W : X
E : F :: *e* : *f*	E : F :: R : S	*e* : *f* :: X : Y
and G : H :: *g* : *h*	G : H :: S : T	*g* : *h* :: Y : Z

then P : T = V : Z.

For $\frac{P}{Q} = \frac{A}{B} = \frac{a}{b} = \frac{V}{W}$,

$\frac{Q}{R} = \frac{C}{D} = \frac{c}{d} = \frac{W}{X}$,

$\frac{R}{S} = \frac{E}{F} = \frac{e}{f} = \frac{X}{Y}$,

$\frac{S}{T} = \frac{G}{H} = \frac{g}{h} = \frac{Y}{Z}$;

and $\therefore \frac{P \times Q \times R \times S}{Q \times R \times S \times T} = \frac{V \times W \times X \times Y}{W \times X \times Y \times Z}$,

and $\therefore \frac{P}{T} = \frac{V}{Z}$,

or P : T = V : Z.

$\therefore$ If ſeveral ratios, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *a ratio which is compounded of ſeveral ratios be the ſame to a ratio which is compounded of ſeveral other ratios; and if one of the firſt ratios, or the ratio which is compounded of ſeveral of them, be the ſame to one of the laſt ratios, or to the ratio which is compounded of ſeveral of them; then the remaining ratio of the firſt, or, if there be more than one, the ratio compounded of the remaining ratios, ſhall be the ſame to the remaining ratio of the laſt, or, if there be more than one, to the ratio compounded of theſe remaining ratios.*

A B C D E F G H
P Q R S T X

Let A : B, B : C, C : D, D : E, E : F, F : G, G : H, be the firſt ratios, and P : Q, Q : R, R : S, S : T, T : X, the other ratios; alſo, let A : H, which is compounded of the firſt ratios, be the ſame as the ratio of P : X, which is the ratio compounded of the other ratios; and, let the ratio of A : E, which is compounded of the ratios of A : B, B : C, C : D, D : E, be the ſame as the ratio of P : R, which is compounded of the ratios P : Q, Q : R.

Then the ratio which is compounded of the remaining firſt ratios, that is, the ratio compounded of the ratios E : F, F : G, G : H, that is, the ratio of E : H, ſhall be the ſame as the ratio of R : X, which is compounded of the ratios of R : S, S : T, T : X, the remaining other ratios.

Harvard University - Houghton Library / pga_typ_805_47_3730

Becauſe $\frac{A \times B \times C \times D \times E \times F \times G}{B \times C \times D \times E \times F \times G \times H} = \frac{P \times Q \times R \times S \times T}{Q \times R \times S \times T \times X}$,

or $\frac{A \times B \times C \times D}{B \times C \times D \times E} \times \frac{E \times F \times G}{F \times G \times H} = \frac{P \times Q}{Q \times R} \times \frac{R \times S \times T}{S \times T \times X}$,

and $\frac{A \times B \times C \times D}{B \times C \times D \times E} = \frac{P \times Q}{Q \times R}$,

$\therefore \frac{E \times F \times G}{F \times G \times H} = \frac{R \times S \times T}{S \times T \times X}$,

$\therefore \frac{E}{H} = \frac{R}{X}$,

$\therefore E : H = R : X$.

$\therefore$ If a ratio which, &c.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *there be any number of ratios, and any number of other ratios, ſuch that the ratio which is compounded of ratios, which are the ſame to the firſt ratios, each to each, is the ſame to the ratio which is compounded of ratios, which are the ſame, each to each, to the laſt ratios—and if one of the firſt ratios, or the ratio which is compounded of ratios, which are the ſame to ſeveral of the firſt ratios, each to each, be the ſame to one of the laſt ratios, or to the ratio which is compounded of ratios, which are the ſame, each to each, to ſeveral of the laſt ratios—then the remaining ratio of the firſt; or, if there be more than one, the ratio which is compounded of ratios, which are the ſame, each to each, to the remaining ratios of the firſt, ſhall be the ſame to the remaining ratio of the laſt; or, if there be more than one, to the ratio which is compounded of ratios, which are the ſame, each to each, to theſe remaining ratios.*

h k m n s
A B, C D, E F, G H, K L, M N, *a b c d e f g*
O P, Q R, S T, V W, X Y, *h k l m n p*
a b c d e f g

Let A : B, C : D, E : F, G : H, K : L, M : N, be the firſt ratios, and O : P, Q : R, S : T, V : W, X : Y, the other ratios;

and let A : B = *a* : *b*,
C : D = *b* : *c*,
E : F = *c* : *d*,
G : H = *d* : *e*,
K : L = *e* : *f*,
M : N = *f* : *g*.

Harvard University - Houghton Library / pga_typ_805_47_3730

Then, by the definition of a compound ratio, the ratio of $a : g$ is compounded of the ratios of $a : b$, $b : c$, $c : d$, $d : e$, $e : f$, $f : g$, which are the ſame as the ratio of A : B, C : D, E : F, G : H, K : L, M : N, each to each.

Alſo, $O : P = h : k$,
$Q : R = k : l$,
$S : T = l : m$,
$V : W = m : n$,
$X : Y = n : p$.

Then will the ratio of $h : p$ be the ratio compounded of the ratios of $h : k$, $k : l$, $l : m$, $m : n$, $n : p$, which are the ſame as the ratios of O : P, Q : R, S : T, V : W, X : Y, each to each.

$\therefore$ by the hypotheſis $a : g = h : p$.

Alſo, let the ratio which is compounded of the ratios of A : B, C : D, two of the firſt ratios (or the ratios of $a : c$, for $A : B = a : b$, and $C : D = b : c$), be the ſame as the ratio of a : d, which is compounded of the ratios of a : b, b : c, c : d, which are the ſame as the ratios of O : P, Q : R, S : T, three of the other ratios.

And let the ratios of h : s, which is compounded of the ratios of h : k, k : m, m : n, n : s, which are the ſame as the remaining firſt ratios, namely, E : F, G : H, K : L, M : N; alſo, let the ratio of e : g, be that which is compounded of the ratios e : f, f : g, which are the ſame, each to each, to the remaining other ratios, namely, V : W, X : Y. Then the ratio of h : s ſhall be the ſame as the ratio of e : g; or h : s = e : g.

For $\frac{A \times C \times E \times G \times K \times M}{B \times D \times F \times H \times L \times N} = \frac{a \times b \times c \times d \times e \times f}{b \times c \times d \times e \times f \times g}$,

Harvard University - Houghton Library / pga_typ_805_47_3730

and $\frac{O \times Q \times S \times V \times X}{P \times R \times T \times W \times Y} = \frac{h \times k \times l \times m \times n}{k \times l \times m \times n \times p}$,

by the compoſition of the ratios;

$\therefore \frac{a \times b \times c \times d \times e \times f}{b \times c \times d \times e \times f \times g} = \frac{h \times k \times l \times m \times n}{k \times l \times m \times n \times p}$ (hyp.),

or $\frac{a \times b}{b \times c} \times \frac{c \times d \times e \times f}{d \times e \times f \times g} = \frac{h \times k \times l}{k \times l \times m} \times \frac{m \times n}{n \times p}$,

but $\frac{a \times b}{b \times c} = \frac{A \times C}{B \times D} = \frac{O \times Q \times S}{P \times R \times T} = \frac{a \times b \times c}{b \times c \times d} = \frac{h \times k \times l}{k \times l \times m}$;

$\therefore \frac{c \times d \times e \times f}{d \times e \times f \times g} = \frac{m \times n}{n \times p}$.

And $\frac{c \times d \times e \times f}{d \times e \times f \times \delta} = \frac{h \times k \times m \times n}{k \times m \times n \times s}$ (hyp.),

and $\frac{m \times n}{n \times p} = \frac{e \times f}{f \times g}$ (hyp.),

$\therefore \frac{h \times k \times m \times n}{k \times m \times n \times s} = \frac{e\,f}{f\,g}$,

$\therefore \frac{h}{s} = \frac{e}{g}$,

$\therefore h : s = e : g.$

$\therefore$ If there be any number, &c.

*** Algebraical and Arithmetical expoſitions of the Fifth Book of Euclid are given in Byrne's Doctrine of Proportion; published by WILLIAMS and Co. London. 1841.

Harvard University - Houghton Library / pga_typ_805_47_3730

BOOK VI.

DEFINITIONS.

I.

RECTILINEAR figures are ſaid to be ſimilar, when they have their ſeveral angles equal, each to each, and the ſides about the equal angles proportional.



II.

Two ſides of one figure are ſaid to be reciprocally proportional to two ſides of another figure when one of the ſides of the firſt is to the ſecond, as the remaining ſide of the ſecond is to the remaining ſide of the firſt.

III.

A STRAIGHT line is ſaid to be cut in extreme and mean ratio, when the whole is to the greater ſegment, as the greater ſegment is to the leſs.


IV.

THE altitude of any figure is the ſtraight line drawn from its vertex perpendicular to its baſe, or the baſe produced.



Harvard University - Houghton Library / pga_typ_805_47_3730

TRIANGLES *and parallelograms having the ſame altitude are to one another as their baſes.*



Let the triangles  and  have a common vertex, and their baſes —— and —— in the ſame ſtraight line.



Produce —— both ways, take ſucceſſively on —— produced lines equal to it; and on —— produced lines ſucceſſively equal to it; and draw lines from the common vertex to their extremities.




The triangles  thus formed are all equal to one another, ſince their baſes are equal. (B. 1. pr. 38.)

∴  and its baſe are reſpectively equi-multiples of  and the baſe ——.

Harvard University - Houghton Library / pga_typ_805_47_3730

In like manner  and its baſe are reſpectively equimultiples of  and the baſe .

∴ If *m* or 6 times   or  *n* or 5 times 

then *m* or 6 times   or  *n* or 5 times  ,


m and *n* ſtand for every multiple taken as in the fifth definition of the Fifth Book. Although we have only ſhown that this property exiſts when *m* equal 6, and *n* equal 5, yet it is evident that the property holds good for every multiple value that may be given to *m*, and to *n*.


∴  : :: : (B. 5. def. 5.)


Parallelograms having the ſame altitude are the doubles of the triangles, on their baſes, and are proportional to them (Part 1), and hence their doubles, the parallelograms, are as their baſes. (B. 5. pr. 15.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730







IF *a ſtraight line* —— *be drawn parallel to any ſide* ▪▪▪▪▪▪▪▪▪▪ *of a triangle, it ſhall cut the other ſides, or thoſe ſides produced, into proportional ſegments.*


And if any ſtraight line —— *divide the ſides of a triangle, or thoſe ſides produced, into proportional ſegments, it is parallel to the remaining ſide* ▪▪▪▪▪▪▪▪▪▪.

PART I.


Let —— ∥ ▪▪▪▪▪▪▪▪, then ſhall

—— : ▪▪▪▪▪▪▪▪ :: ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪.

Draw —— and ——,

and  (B. 1. pr. 37);

 (B.5. pr.7); but

 (B. 6. pr. 1),


∴ —— : ▪▪▪▪▪▪▪▪ :: ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪.

(B. 5. pr. 11).


Harvard University - Houghton Library / pga_typ_805_47_3730

PART II.

Let  ,


then  .


Let the ſame conſtruction remain,


becauſe  (B. 6. pr. 1);

and

but  (hyp.),


∴  (B. 5. pr. 11.)

∴  (B. 5. pr. 9);

but they are on the ſame baſe  , and at the ſame ſide of it, and

∴  (B. 1. pr. 39).

Q. E. D.



A RIGHT *line* (——) *biſecting the angle of a triangle, divides the oppoſite ſide into ſegments* (——, - - - - - - -) *proportional to the conterminous ſides* (——, ——).

And if a ſtraight line (——) *drawn from any angle of a triangle divide the oppoſite ſide* (——————) *into ſegments* (——, ——————) *proportional to the conterminous ſides* (——, ——), *it biſects the angle.*

PART I.

Draw - - - - - - - - || ——, to meet - - - - - - - - - -;

then, ◢ = ◢ (B. 1. pr. 29),

∴ ◣ = ◢; but ◣ = ◥, ∴ ◥ = ◢,

∴ - - - - - - - - - - = —— (B. 1. pr. 6);

and becauſe —— || - - - - - - - - - -,

- - - - - - - - - - : —— :: - - - - - - - - : ——

(B. 6. pr. 2);

but - - - - - - - - - - = ——;


∴ —— : —— :: - - - - - - - - : ——


(B. 5. pr. 7).


Harvard University - Houghton Library / pga_typ_805_47_3730

PART II.


Let the ſame conſtruction remain,

and  (B. 6. pr. 2);

but  (hyp.)


∴ 


(B. 5. pr. 11).


and ∴  (B. 5. pr. 9),

and ∴  (B. 1. pr. 5); but ſince


 ;

and  (B. 1. pr. 29);

∴  , and  ,

and ∴  bifects .

Q. E. D.



IN *equiangular triangles* (and) *the ſides about the equal angles are proportional, and the ſides which are oppoſite to the equal angles are homologous.*

Let the equiangular triangles be ſo placed that two ſides ——, - - - - - - oppoſite to equal angles and may be conterminous, and in the ſame ſtraight line; and that the triangles lying at the ſame ſide of that ſtraight line, may have the equal angles not conterminous,

i. e. oppoſite to , and to .

Draw and . Then, becauſe = , || (B. 1. pr. 28);

and for a like reaſon, || ,

∴ is a parallelogram.

But : :: : (B. 6. pr. 2);

Harvard University - Houghton Library / pga_typ_805_47_3730

and ſince —— = —— (B. 1. pr. 34),

—— : ····· :: —— : ····· ; and by

alternation, —— : —— :: ····· : ·····

(B. 5. pr. 16).

In like manner it may be ſhown, that

—— : ····· :: —— : ····· ;

and by alternation, that

—— : —— :: ····· : ····· ;

but it has been already proved that

—— : —— :: ····· : ····· ,

and therefore, ex æquali,


—— : —— :: ····· : ·····

(B. 5. pr. 22),

therefore the ſides about the equal angles are proportional, and thoſe which are oppoſite to the equal angles are homologous.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *two triangles have their ſides proportional* (▬▬ : ▬▬ :: ▬▬ : ▬▬) and (▬▬ : ▬▬ :: ▬▬ : ▬▬) *they are equiangular, and the equal angles are ſubtended by the homologous ſides.*

From the extremities of ▬▬, draw ▬▬ and ▬▬, making

▼ = ▲,

● = ◗ (B. 1. pr. 23);

and conſequently ◤ = ◢ (B. 1. pr. 32),

and ſince the triangles are equiangular,

▬▬ : ▬▬ :: ▬▬ : ▬▬

(B. 6. pr. 4);

but ▬▬ : ▬▬ :: ▬▬ : ▬▬ (hyp.);

∴ ▬▬ : ▬▬ :: ▬▬ : ▬▬,


and conſequently ▬▬ = ▬▬ (B. 5. pr. 9).


In the like manner it may be ſhown that


▬▬ = ▬▬.


Harvard University - Houghton Library / pga_typ_805_47_3730


Therefore, the two triangles having a common baſe ———, and their ſides equal, have alſo equal angles oppoſite to equal ſides, i. e.

 (B. 1. pr. 8).

But  (conſt.)

and ∴  ; for the ſame


reaſon  , and

conſequently  (B. 1. 32);

and therefore the triangles are equiangular, and it is evident that the homologous ſides ſubtend the equal angles.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *two triangles* (△ and △) *have one angle* (◔) *of the one, equal to one angle* (◔) *of the other, and the ſides about the equal angles proportional, the triangles ſhall be equiangular, and have thoſe angles equal which the homologous ſides ſubtend.*

From the extremities of ——, one of the ſides of △, about ◔, draw —— and ┈┈, making ◥ = ◔, and ◢ = ◢; then ◥ = ◥

(B. 1. pr. 32), and two triangles being equiangular,

┈┈ : ┈┈┈ :: ┈┈ : ——

(B. 6. pr. 4);

but ┈┈ : ┈┈┈ :: —— : —— (hyp.);

∴ ┈┈ : —— :: —— : ——


(B. 5. pr. 11),


and conſequently ┈┈ = —— (B. 5. pr. 9);


Harvard University - Houghton Library / pga_typ_805_47_3730


$\therefore$  in every reſpect.

(B. 1. pr. 4).

But  (conſt.),

and $\therefore$  ; and


ſince alſo  ,

 (B. 1. pr. 32);

and $\therefore$  and  are equiangular, with their equal angles oppoſite to homologous ſides.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *two triangles* (and) *have one angle in each equal* (*equal to*), *the ſides about two other angles proportional* (: :: :), *and each of the remaining angles* (*and*) *either leſs or not leſs than a right angle, the triangles are equiangular, and thoſe angles are equal about which the ſides are proportional.*

Firſt let it be aſſumed that the angles and are each leſs than a right angle: then if it be ſuppoſed that and contained by the proportional ſides, are not equal, let be the greater, and make

$$= .$$

Becauſe = (hyp.), and = (conſt.)

$$\therefore \quad = \quad \text{(B. 1. pr. 32);}$$

Harvard University - Houghton Library / pga_typ_805_47_3730

∴ —— : —— :: ····· : ·····

(B. 6. pr. 4),

but —— : —— :: ····· : ····· (hyp.)

∴ —— : —— :: —— : ——;

∴ —— = —— (B. 5. pr. 9),

and ∴  =  (B. 1. pr. 5).

But  is leſs than a right angle (hyp.)

∴  is leſs than a right angle; and ∴  muſt be greater than a right angle (B. 1. pr. 13), but it has been proved =  and therefore leſs than a right angle, which is abſurd. ∴  and  are not unequal;

∴ they are equal, and ſince  =  (hyp.)

∴  =  (B. 1. pr. 32), and therefore the triangles are equiangular.

But if  and  be aſſumed to be each not leſs than a right angle, it may be proved as before, that the triangles are equiangular, and have the ſides about the equal angles proportional. (B. 6. pr. 4).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IN *a right angled triangle* ( ), *if a perpendicular* ( ) *be drawn from the right angle to the oppoſite ſide, the triangles* (,) *on each ſide of it are ſimilar to the whole triangle and to each other.*

Becauſe = (B. 1. ax. 11), and

common to and ;

= (B. 1. pr. 32);


∴ and are equiangular; and conſequently have their ſides about the equal angles proportional (B. 6. pr. 4), and are therefore ſimilar (B. 6. def. 1).

In like manner it may be proved that is ſimilar to ; but has been ſhewn to be ſimilar to ; ∴ and are ſimilar to the whole and to each other.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

FROM *a given ſtraight line* (——) *to cut off any required part.*



From either extremity of the given line draw —— making any angle with ——; and produce —— till the whole produced line —— contains —— as often as —— contains the required part.

Draw ——, and draw —— ‖ ——.

—— is the required part of ——.

For ſince —— ‖ ——

—— : —— :: —— : ——

(B. 6. pr. 2), and by compoſition (B. 5. pr. 18);


—— : —— :: —— : ——;

but —— contains —— as often as —— contains the required part (conſt.);

∴ —— is the required part.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



TO *divide a ſtraight line* (——) *ſimilarly to a given divided line* (——).

From either extremity of the given line —— draw ——— making any angle; take ———, ——— and ——— equal to ——, ——— and ——— reſpectively (B. 1. pr. 2); draw ———, and draw ——— and ——— ‖ to it.

Since { ——— } are ‖,

——— : ——— :: ——— : ———

(B. 6. pr. 2),

or ——— : ——— :: ——— : ——— (conſt.),

and ——— : ——— :: ——— : ———

(B. 6. pr. 2),

——— : ——— :: ——— : ——— (conſt.),

and ∴ the given line —— is divided ſimilarly to ——.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *find a third proportional to two given ſtraight lines* (——— *and* ———).



At either extremity of the given line ——— draw ----- making an angle; take ---------- = ———, and draw ———— ; make ---------- = ———, and draw ---------- || ———— ; (B. 1. pr. 31.) ——— is the third proportional to ——— and ———.

For ſince ———— || ----------,

∴ ——— : ---------- :: ---------- : ———

(B. 6 pr. 2);

but ---------- = ---------- = ——— (conſt.);

∴ ——— : ——— :: ——— : ———.

(B. 5. pr. 7).

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730



TO *find a fourth proportional to three given lines*

{ — — —

TO *find a mean proportional between two given ſtraight lines* { ——— } .



Draw any ſtraight line ——— ,

make ——— = ——— ,

and ——— = ——— ; biſect ——— : and from the point of biſection as a centre, and half the line as a radius, deſcribe a ſemicircle ⌒ ,

draw ——— ⊥ ——— :

——— is the mean proportional required.

Draw ——— and ——— .

Since ◔ is a right angle (B. 3. pr. 31),

and ——— is ⊥ from it upon the oppoſite ſide,

∴ ——— is a mean proportional between ——— and ——— (B. 6. pr. 8),

and ∴ between ——— and ——— (conſt.).

Q. E. D

Harvard University - Houghton Library / pga_typ_805_47_3730



I.

EQUAL *parallelograms*  *and*  , *which have one angle in each equal, have the ſides about the equal angles reciprocally proportional* (— : — :: — : —).

II.

And parallelograms which have one angle in each equal, and the ſides about them reciprocally proportional, are equal.

Let — and —; and — and —, be ſo placed that — and — may be continued right lines. It is evident that they may aſſume this poſition. (B. 1. prs. 13, 14, 15.)

Complete  .

Since  ;

∴  (B. 5. pr. 7.)

∴ — : — :: — : —

(B. 6. pr. 1.)

The ſame conſtruction remaining:

— : — :: { ▱ : ▱ (B. 6. pr. 1.)
{ — : — (hyp.)

▱ : ▱ (B. 6. pr. 1.)

∴ ▱ : ▱ :: ▱ : ▱ (B. 5. pr. 11.)

and ∴ ▱ = ▱ (B. 5. pr. 9).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



I.

EQUAL *triangles, which have one angle in each equal* (=), *have the ſides about the equal angles reciprocally proportional* (: :: :).

II.

And two triangles which have an angle of the one equal to an angle of the other, and the ſides about the equal angles reciprocally proportional, are equal.

I.

Let the triangles be ſo placed that the equal angles and may be vertically oppoſite, that is to ſay, ſo that and may be in the ſame ſtraight line. Whence alſo and muſt be in the ſame ſtraight line. (B. 1. pr. 14.)

Draw , then

: :: : (B. 6. pr. 1.)

:: : (B. 5. pr. 7.)

:: : (B. 6. pr. 1.)

Harvard University - Houghton Library / pga_typ_805_47_3730

∴ —— : —— :: —— : ——

(B. 5. pr. 11.)

II.

Let the ſame conſtruction remain, and

▶ : ▼ :: —— : —— (B. 6. pr. 1.)

and —— : —— :: ◀ : ▼

(B. 6. pr. 1.)

But —— : —— :: —— : ——, (hyp.)

∴ ▶ : ▼ :: ◀ : ▼ (B. 5 pr. 11);

∴ ▶ = ◀ (B. 5. pr. 9.)

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



PART I.

IF *four ſtraight lines be proportional* (—— : —— :: ---------- : ---------), *the rectangle* (—— × ---------) *contained by the extremes, is equal to the rectangle* (—— × ----------) *contained by the means.*



PART II.

And if the rectangle contained by the extremes be equal to the rectangle contained by the means, the four ſtraight lines are proportional.

PART I.

From the extremities of —— and —— draw —— and —— ⊥ to them and = ---------- and ---------- reſpectively: complete the parallelograms  and  .

And ſince,

—— : —— :: ---------- : ---------- (hyp.)

∴ —— : —— :: —— : —— (conſt.)


∴  =  (B. 6. pr. 14),

Harvard University - Houghton Library / pga_typ_805_47_3730

that is, the rectangle contained by the extremes, equal to the rectangle contained by the means.

PART II.

Let the ſame conſtruction remain; becauſe






(B. 6. pr. 14).



(B. 5. pr. 7).

Q. E. D.



PART I

IF *three ſtraight lines be proportional* (— : — :: — : —) *the rectangle under the extremes is equal to the ſquare of the mean.*

PART II.

And if the rectangle under the extremes be equal to the ſquare of the mean, the three ſtraight lines are proportional.

PART I.

Aſſume — = —, and

ſince — : — :: — : —,

then — : — :: — : —,

∴ — × — = — × —

(B. 6. pr. 16).

But — = —,

∴ — × — = — × —,

or = —^{2}; therefore, if the three ſtraight lines are proportional, the rectangle contained by the extremes is equal to the ſquare of the mean.

PART II.

Aſſume — = —, then

— × — = — × —,

∴ — : — :: — : —

(B. 6. pr. 16), and

∴ — : — :: — : —.

Q. E. D.


Harvard University - Houghton Library / pga_typ_805_47_3730


ON *a given ſtraight line* (——) *to conſtruct a rectilinear figure*  *ſimilar to a given one* ( ) *and ſimilarly placed.*





Reſolve the given figure into triangles by drawing the lines - - - - - - - - and . . . . . .



At the extremities of —— make  and  :

again at the extremities of —— make 

and  : in like manner make

 .

Then  is ſimilar to  .

It is evident from the conſtruction and (B. 1. pr. 32) that the figures are equiangular; and ſince the triangles

 and  are equiangular; then by (B. 6. pr. 4),


——— : —— :: - - - - - - - - - : ——

and —— : —— :: —— : . . . . . .

Again, becauſe  and  are equiangular,

 ;

∴ ex æquali,



(B. 6. pr. 22.)

In like manner it may be ſhown that the remaining ſides of the two figures are proportional.

∴ by (B. 6. def. 1.)

 is ſimilar to 


and ſimilarly ſituated; and on the given line .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

SIMILAR *trian-*

gles ( *and* ) *are to one another in the duplicate ratio of their homologous ſides.*



Let  and  be equal angles, and ▪▪▪▪▪▪▪▪ and ▪▪▪▪▪▪▪▪ homologous ſides of the ſimilar triangles  and  and on ▪▪▪▪▪▪▪▪ the greater of theſe lines take ▪▪▪▪▪▪▪ a third proportional, ſo that

▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪ :: ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪▪▪;

draw ▪▪▪▪▪▪▪▪▪▪.

▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪ :: ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪


(B. 6. pr. 4);

∴ ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪ :: ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪

(B. 5. pr. 16, alt.),


but ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪ :: ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪ (conſt.),

∴ ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪ :: ▪▪▪▪▪▪▪▪ : ▪▪▪▪▪▪▪▪ conſe-

quently  for they have the ſides about

the equal angles  and  reciprocally proportional

(B. 6. pr. 15);




(B. 5 pr. 7);

but 

(B. 6. pr. 1),




that is to ſay, the triangles are to one another in the duplicate ratio of their homologous ſides


 and  (B. 5. def. 11).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



SIMILAR *polygons may be divided into the ſame number of ſimilar triangles, each ſimilar pair of which are proportional to the polygons; and the polygons are to each other in the duplicate ratio of their homologous ſides.*



Draw ——— and ▪▪▪▪▪▪▪▪▪▪▪, and ——— and ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪, reſolving the polygons into triangles. Then becauſe the polygons are ſimilar, ◢ = ◢,

and ——— : ▪▪▪▪▪▪▪▪▪▪▪ :: ——— : ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪

∴ ◢ and ◢ are ſimilar, and ◀ = ◀

(B. 6. pr. 6);

but ◥ = ◥ becauſe they are angles of ſimilar polygons; therefore the remainders ▲ and ▲ are equal;

hence ▪▪▪▪▪▪▪▪▪▪▪ : ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ :: ▪▪▪▪▪▪▪▪▪ : ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪ ▪

and ---------- : ——— :: ---------- : ———,

on account of the ſimilar polygons,

∴ ---------- : ——— :: ---------- : ———,

ex æquali (B. 5. pr. 22), and as theſe proportional ſides contain equal angles, the triangles  and  are ſimilar (B. 6. pr. 6).

In like manner it may be ſhown that the triangles  and  are ſimilar.


But  is to  in the duplicate ratio of ---------- to ---------- (B. 6. pr. 19), and  is to  in like manner, in the duplicate ratio of ---------- to ----------;

∴  :  ::  :  ,

(B. 5. pr. 11);


Again  is to  in the duplicate ratio of ——— to ———, and  is to  in

the duplicate ratio of ——— to ———,



and as one of the antecedents is to one of the conſequents, ſo is the ſum of all the antecedents to the ſum of all the conſequents; that is to ſay, the ſimilar triangles have to one another the ſame ratio as the polygons (B. 5. pr. 12).

But  is to in the duplicate ratio of ——— to ———;



∴ is to in the duplicate ratio of ——— to ———.

Q. E. D



RECTILINEAR *figures* ( *and* ) *which are ſimilar to the ſame figure* () *are ſimilar alſo to each other.*





Since  and  are ſimilar, they are equiangular, and have the ſides about the equal angles proportional (B. 6. def. 1); and ſince the figures  and  are alſo ſimilar, they are equiangular, and have the ſides about the equal angles proportional; therefore  and  are alſo equiangular, and have the ſides about the equal angles proportional (B. 5. pr. 11), and are therefore ſimilar.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730


PART I.

IF *four ſtraight lines be proportional* (— : — :: — : —), *the ſimilar rectilinear figures ſimilarly described on them are alſo proportional.*



PART II.

And if four ſimilar rectilinear figures, ſimilarly deſcribed on four ſtraight lines, be proportional, the ſtraight lines are alſo proportional.



PART I.

Take ····· a third proportional to —

and ——, and ····· a third proportional

to —— and —— (B. 6. pr. 11);

ſince — : —— :: —— : —— (hyp.),

—— : ····· :: —— : ····· (conſt.)

∴ ex æquali,

— : —— :: —— : ·····;

but  :  :: — : ·····

(B. 6. pr. 20),


and  :  :: —— : ·····;




(B. 5. pr. 11).

PART II.

Let the ſame conſtruction remain:


(hyp.),


(conſt.)

and
.

(B. 5. pr. 11).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

EQUIANGULAR *parallelograms* (and) *are to one another in a ratio compounded of the ratios of their ſides.*



Let two of the ſides —— and ..... about the equal angles be placed ſo that they may form one ſtraight line.

Since + = ,

and = (hyp.),

+ = ,

and ∴ —— and —— form one ſtraight line (B. 1. pr. 14);

complete .

Since : :: —— : .....


(B. 6. pr. 1),

and : :: —— : —— (B. 6. pr. 1),

has to a ratio compounded of the ratios of —— to ....., and of —— to ——.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IN any parallelogram ( ) the parallelograms (and) which are about the diagonal are ſimilar to the whole, and to each other.

As and have a common angle they are equiangular; but becauſe ||

and are ſimilar (B. 6. pr. 4),

∴ : :: : ;

and the remaining oppoſite ſides are equal to thoſe,

∴ and have the ſides about the equal angles proportional, and are therefore ſimilar.

In the ſame manner it can be demonſtrated that the parallelograms and are ſimilar.

Since, therefore, each of the parallelograms and is ſimilar to , they are ſimilar to each other.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *deſcribe a rectilinear figure, which ſhall be ſimilar to a given rectilinear figure* ( ), *and equal to another* ( ).



Upon —— deſcribe ■ = ◺,

and upon —— deſcribe □ = ⬠,

and having ◔ = ◔ (B. 1. pr. 45), and then

—— and ┅┅┅ will lie in the ſame ſtraight line

(B. 1. prs. 29, 14),

Between —— and ┅┅┅ find a mean proportional

—— (B. 6. pr. 13), and upon ——

deſcribe ◺, ſimilar to ◺,

and ſimilarly ſituated.

Then ◺ = ⬠.

For ſince ◺ and ◺ are ſimilar, and

—— : —— :: —— : ┅┅┅ (conſt.),

◺ : ◺ :: —— : ┅┅┅

(B. 6. pr. 20);

Harvard University - Houghton Library / pga_typ_805_47_3730

but  :  :: :  (B. 6. pr. 1);

$\therefore$  :  ::  :  (B. 5. pr. 11);

but  =  (conſt.),

and $\therefore$  =  (B. 5. pr. 14);


and  =  (conſt.); conſequently,

 which is ſimilar to  is alſo =  .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *similar and similarly posited parallelograms* (and) *have a common angle, they are about the same diagonal.*



For, if possible, let be the diagonal of and draw ‖ (B. 1. pr. 31).

Since and are about the same diagonal , and have common, they are similar (B. 6. pr. 24);

∴ : :: : ;

but : :: :

(hyp.),

∴ : :: : ,

and ∴ = (B. 5. pr. 9.),

which is absurd.

∴ is not the diagonal of



in the same manner it can be demonstrated that no other line is except .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



OF *all the rectangles contained by the ſegments of a given ſtraight line, the greateſt is the ſquare which is deſcribed on half the line.*

Let be the given line,  and  unequal ſegments, and and equal ſegments;


then  .

For it has been demonſtrated already (B. 2. pr. 5), that the ſquare of half the line is equal to the rectangle contained by any unequal ſegments together with the ſquare of the part intermediate between the middle point and the point of unequal ſection. The ſquare deſcribed on half the line exceeds therefore the rectangle contained by any unequal ſegments of the line.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *divide a given ſtraight line* ( ) *ſo that the rectangle contained by its segments may be equal to a given area, not exceeding the ſquare of half the line.*



Let the given area be $=$ ^{2}.

Biſect , or

make $=$;

and if $^2 =$ ^{2},

the problem is ſolved.

But if $^2 \neq$ ^{2}, then

muſt $\sqsubset$ (hyp.).

Draw $\perp$ $=$;

make $=$ or ;

with as radius deſcribe a circle cutting the given line; draw .

Then $\times$ $+$ $^2 =$ ^{2}

(B. 2. pr. 5.) $=$ ^{2}.

But $^2 =$ $^2 +$ ^{2}

(B. 1. pr. 47);

Harvard University - Houghton Library / pga_typ_805_47_3730

$\therefore$ ⟋ × ⟋ + ⟋^{2}

$= ⟋^2 + ⟋^2$,

from both, take $⟋^2$,

and ⟋ × ⟋ $= ⟋^2$.

But ⟋ = ⟋ (conſt.),


and $\therefore$ ⟋ is ſo divided

that ⟋ × ⟋ $= ⟋^2$.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

TO *produce a given ſtraight line* (———), *ſo that the rectangle contained by the ſegments between the extremities of the given line and the point to which it is produced, may be equal to a given area, i. e. equal to the ſquare on* ———.



Make ——— = ———, and
draw ——— ⊥ ——— = ———;
draw ———; and
with the radius ———, deſcribe a circle
meeting ——— produced.


Then ——— × ——— + ———^{2} =
———^{2} (B. 2. pr. 6.) = ———^{2}.

But ———^{2} = ———^{2} + ———^{2} (B. 1. pr. 47.)

∴ ——— × ——— + ———^{2} =
———^{2} + ———^{2},
from both take ———^{2},
and ——— × ——— = ———^{2};
but ——— = ———,

∴ ———^{2} = the given area.

Q. E. D.



TO *cut a given finite ſtraight line* (——·····) *in extreme and mean ratio.*

On ——····· deſcribe the ſquare 

(B. 1. pr. 46); and produce ———, ſo that

——···· × ····· = ——·····^{2}

(B. 6. pr. 29);

take —— = ·····,

and draw ——— ‖ ——····,

meeting —— ‖ ——····· (B. 1. pr. 31).

Then  = —— × ·····,

and is ∴ =  ; and if from both theſe equals

be taken the common part  ,

 , which is the ſquare of ——,

will be =  , which is = ——— × ·····;

that is ——^{2} = ——···· × ·····;

∴ ——····· : —— :: —— : ·····,


and ——····· is divided in extreme and mean ratio.

(B. 6. def. 3).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *any similar rectilinear figures be similarly described on the sides of a right angled triangle* (), *the figure described on the side* ( ) *subtending the right angle is equal to the sum of the figures on the other sides.*



From the right angle draw ——— perpendicular to ——— ;

then ——— : ——— :: ——— : ———

(B. 6. pr. 8).

∴ ■ : ■ :: ——— : ———

(B. 6. pr. 20).

but ■ : ■ :: ——— : ———

(B. 6. pr. 20).

Hence ——— + ——— : ———

:: ■ + ■ : ■ ;

but ——— + ——— = ——— ;

and ∴ ■ + ■ = ■ .

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *two triangles* (and), *have two ſides proportional* (: :: :), *and be ſo placed at an angle that the homologous ſides are parallel, the remaining ſides* (*and*) *form one right line.*

Since ||,

= (B. 1. pr. 29);

and alſo ſince ||,

= (B. 1. pr. 29);

∴ = ; and ſince

: :: : (hyp.),

the triangles are equiangular (B. 6. pr. 6);

∴ = :

but = ;

∴ + + = + + =


(B. 1. pr. 32), and ∴ and

lie in the ſame ſtraight line (B. 1. pr. 14).


Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IN *equal circles* (○ , ○), *angles, whether at the centre or circumference, are in the ſame ratio to one another as the arcs on which they ſtand* (◺ : ◺ :: ▬ : ⌒); *ſo alſo are ſectors.*



Take in the circumference of ○ any number of arcs ⌒ , ⌒ , &c. each = ⌒ , and alſo in the circumference of ○ take any number of arcs ⌒ , ⌒ , &c. each = ⌒ , draw the radii to the extremities of the equal arcs.



Then ſince the arcs ⌒ , ⌒ , ⌒ , &c. are all equal, the angles ◺ , ◺ , ◺ , &c. are alſo equal (B. 3. pr. 27);



∴ ◭ is the ſame multiple of ◺ which the arc ⌒ is of ▬ ; and in the ſame manner ◭ is the ſame multiple of ◺ , which the arc ⌒ is of the arc ⌒ .



Harvard University - Houghton Library / pga_typ_805_47_3730

Then it is evident (B. 3. pr. 27),

if  (or if m times ) $\sqsubset, =, \sqsupset$ 

(or n times )

then  (or m times ) $\sqsubset, =, \sqsupset$

 (or n times );

$\therefore$  , (B. 5. def. 5), or the angles at the centre are as the arcs on which they ſtand; but the angles at the circumference being halves of the angles at the centre (B. 3. pr. 20) are in the ſame ratio (B. 5. pr. 15), and therefore are as the arcs on which they ſtand.


It is evident, that ſectors in equal circles, and on equal arcs are equal (B. 1. pr. 4; B. 3. prs. 24, 27, and def. 9). Hence, if the ſectors be ſubſtituted for the angles in the above demonſtration, the ſecond part of the propoſition will be eſtabliſhed, that is, in equal circles the ſectors have the ſame ratio to one another as the arcs on which they ſtand.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730

IF *the right line* (·····), *biſecting an external angle*  *of the tri-angle*  *meet the oppoſite*  *ſide* (——) *produced, that whole produced ſide* (——·····), *and its external ſegment* (·····) *will be proportional to the ſides* (—····· *and* ——), *which contain the angle adjacent to the external biſected angle.*

For if —— be drawn ‖ ·····,

then  = , (B. 1. pr. 29);

=  , (hyp.),

=  , (B. 1. pr. 29);

and ∴ ····· = ——, (B. 1. pr. 6),

and —····· : —— :: —····· : ·····,

(B. 5. pr. 7);

But alſo,

——····· : ····· :: —····· : ·····,

(B. 6. pr. 2);


and therefore

——····· : ····· :: —····· : ——,

(B. 5. pr. 11).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *an angle of a triangle be bi-ſected by a ſtraight line, which likewiſe cuts the baſe; the rectangle contained by the ſides of the triangle is equal to the rectangle contained by the ſegments of the baſe, together with the ſquare of the ſtraight line which biſects the angle.*

Let —— be drawn, making ◢ = ◣; then ſhall

—— × —— = ····· × —— + ——².

About ◬ deſcribe ○ (B. 4. pr. 5),

produce —— to meet the circle, and draw ·····.

Since ◢ = ◭ (hyp.),

and ◿ = ◥ (B. 3. pr. 21),

∴ ◬ and ◬ are equiangular (B. 1. pr. 32);

∴ —— : —— :: ····· : ——

(B. 6. pr. 4);

Harvard University - Houghton Library / pga_typ_805_47_3730

$\therefore$ —— × —— = —— × ——

(B. 6. pr. 16.)

= ------ × —— + ——^{2}

(B. 2. pr. 3);


but ------ × —— = ------ × ——

(B. 3. pr. 35);

$\therefore$ —— × —— = ------ × —— + ——^{2}.

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730



IF *from any angle of a triangle a ſtraight line be drawn perpendicular to the baſe; the rectangle contained by the ſides of the triangle is equal to the rectangle contained by the perpendicular and the diameter of the circle deſcribed about the triangle.*

From ▲ of ◺ draw ┈┈ ⊥ ┈┈; then ſhall ┈┈ × ── = ┈┈ × the diameter of the deſcribed circle.

Deſcribe ◯ (B. 4. pr. 5), draw its diameter ──, and draw ──; then becauſe ◔ = ◺ (conſt. and B. 3. pr. 31);

and △ = △ (B. 3. pr. 21);

∴ ◺ is equiangular to △ (B. 6. pr. 4);

∴ ┈┈ : ┈┈ :: ── : ──;


and ∴ ┈┈ × ── = ┈┈ × ──

(B. 6. pr. 16).

Q. E. D.

Harvard University - Houghton Library / pga_typ_805_47_3730


THE *rectangle contained by the diagonals of a quadrilateral figure inſcribed in a circle, is equal to both the rectangles contained by its oppoſite ſides.*




Let  be any quadrilateral figure inſcribed in  ; and draw and ; then




Make  (B. 1. pr. 23),


$\therefore$  ; and (B. 3. pr. 21);

$\therefore$  (B. 6. pr. 4);

and $\therefore$  (B. 6. pr. 16); again,

because  (conſt.),

Harvard University - Houghton Library / pga_typ_805_47_3730

and  (B. 3. pr. 21);




(B. 6. pr. 4);

and 

(B. 6. pr. 16);

but, from above,





(B. 2. pr. 1.

Q. E. D.

THE END.


CHISWICK: PRINTED BY C. WHITTINGHAM.

Harvard University - Houghton Library / pga_typ_805_47_3730

Harvard University - Houghton Library / pga_typ_805_47_3730

Typ 805.47.3730
HOUGHTON LIBRARY
HARVARD COLLEGE LIBRARY

Harvard University - Houghton Library / pga_typ_805_47_3730



Harvard University - Houghton Library / pga_typ_805_47_3730